



## Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Artikel 4  
Assignment title: Henik Sukorini  
Submission title: Biodiesel Industrial Waste based on *Jatropha curcas* as a Fungi...  
File name: Biodiesel Industrial Waste based on *Jatropha curcas* as a Fungi...  
File size: 6.05M  
Page count: 7  
Word count: 3,677  
Character count: 18,065  
Submission date: 06-Aug-2025 10:18AM (UTC+0700)  
Submission ID: 2725899653

E3S Web of Conferences 374, 00016 (2023) <https://doi.org/10.1051/e3sconf/202337400016>  
3<sup>rd</sup> NRLS

---

**Biodiesel Industrial Waste based on *Jatropha curcas* as a Fungicide to Control *Fusarium oxysporum* and *Alternaria solani***

Henik Sukorini<sup>1</sup>, Diah Eri Widyasari<sup>2</sup>, Dini Kurniawati<sup>3</sup>, Sorvin Susannara<sup>4</sup>, Muzivran Mel<sup>5</sup>, and Roy Hendroko Setyobudi<sup>6</sup>

<sup>1</sup>Department of Agrotechnology, Faculty of Agriculture and Animal Science, University of Muhammadiyah Malang, Jl. Raya Tlogomas No. 246, Malang 65144, East Java, Indonesia  
<sup>2</sup>Department of Mechanical Engineering, Faculty of Engineering, University of Muhammadiyah Malang, Malang, 65144, East Java, Indonesia  
<sup>3</sup>Department of Plant Pathology, Faculty of Agriculture, Kasetsart University, 50 Ngamwongwan Rd, Lat Yao, Chachoabuk, Bangkok 10900, Thailand  
<sup>4</sup>Department of Chemical Engineering and Sustainability, Faculty of Engineering, International Islamic University Malaysia, 50728 Kuala Lumpur, Malaysia  
<sup>5</sup>Postgraduate School, Program Study of Biology, Manara Universitas Nasional, Jl. RM. Harsono No.1, Special Region of Jakarta 12550, Indonesia  
<sup>6</sup>Department of Agriculture Science, Postgraduate Program, University of Muhammadiyah Malang, Malang, 65144, East Java, Indonesia

**Abstract.** *Fusarium oxysporum* (Sclerotium, anamniot, Sordari & Hansen.) is the most devastating pathogen causing wilt disease on the tomatoes (*Solanum lycopersicum* L.) plant, whereas *Alternaria solani* (Sordari) is a pathogen that caused early blight on potatoes (*Solanum tuberosum* L.). *Jatropha curcas* L. is a biodiesel material known as a potential fungicide. The industrial biodiesel waste based on *J. curcas* was not yet observed, particularly on the utilization of waste mainly for green manure and biogas. This research aimed to evaluate the extract of industrial biodiesel waste based on *J. curcas* against *F. oxysporum* and *A. solani*. There were 2 × 10 mg L<sup>-1</sup>, 4 × 10 mg L<sup>-1</sup>, 6 × 10 mg L<sup>-1</sup>, 8 × 10 mg L<sup>-1</sup>, and 10 × 10 mg L<sup>-1</sup> of crude extract that were tested on both pathogen *in vitro* and *in vivo*. The concentration of the extract was in an effective range of 10 × 10 mg L<sup>-1</sup> and was able to inhibit the growth of all isolates of *F. oxysporum* from both locations by more than 80 %, but the inhibition of the pathogen *A. solani* was less than 70 %.

**Keywords:** Environmentally friendly, pathogen control, waste to fungicide, waste utilization

**1. Introduction**

*Jatropha curcas* L. is a bioenergy resource in the family Euphorbiaceae. It is a perennial bush that is easy to grow in tropical and subtropical regions and resistant to drought. This plant is native to Mexico and Central America but can be cultivated in South East Asia [1, 2]. Indonesia has started to develop *J. curcas*-based energy to reduce dependence on fossil energy. It is possible because *J. curcas* can be grown on marginal lands that do not compete with food crops [3–6].

Tomato (*Solanum lycopersicum* L.) plants often experience decreased production due mainly to *Fusarium* wilt disease. *Fusarium* attacks tomatoes with a yield reduction of up to 85 % [1]. Until now, this disease is very difficult to control even with synthetic pesticides. Various ways have been done including using antagonistic microorganisms, but the results have not been able to effectively control this disease. The other disease is early blight. Early blight is one of the important diseases in potato plants after *Phytophthora*. The attack rate of this disease reaches 85 % [4].

In the processing of *J. curcas* as a biofuel, the process generates waste, such as seed coat waste, liquid waste, and husk. *J. curcas* seed cake is generated in considerable quantities as a by-product of *J. curcas* seed oil extraction [7, 8]. This biodiesel waste contains toxic ingredients so it cannot be used immediately. Generally, the collection and disposal

\*Corresponding author: [haskornin@yahoo.com](mailto:haskornin@yahoo.com)


© The Authors, published by EDP Sciences. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (<http://creativecommons.org/licenses/by/4.0/>).

## Artikel 4

# Biodiesel Industrial Waste based on *Jatropha curcas* as a Fungicide to Control *Fusarium oxysporum* and *Alternaria solani*

 Henik Sukorini

 Publication Articles Jul - Aug 2025 Dosen UMM

 University of Muhammadiyah Malang

---

### Document Details

Submission ID

trn:oid::1:3308961887

Submission Date

Aug 6, 2025, 10:18 AM GMT+7

Download Date

Aug 6, 2025, 11:40 AM GMT+7

File Name

Biodiesel Industrial Waste based on *Jatropha curcas* as a Fungicide to Control *Fusarium oxyspor*....pdf

File Size

6.1 MB

7 Pages

3,677 Words

18,065 Characters

# 17% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

## Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text
- ▶ Cited Text

## Exclusions

- ▶ 10 Excluded Sources

## Match Groups

- **36 Not Cited or Quoted 17%**  
 Matches with neither in-text citation nor quotation marks
- **0 Missing Quotations 0%**  
 Matches that are still very similar to source material
- **0 Missing Citation 0%**  
 Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted 0%**  
 Matches with in-text citation present, but no quotation marks

## Top Sources

- 16% Internet sources
- 14% Publications
- 8% Submitted works (Student Papers)

## Integrity Flags

### 0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

### Match Groups

- **36 Not Cited or Quoted 17%**  
Matches with neither in-text citation nor quotation marks
- **0 Missing Quotations 0%**  
Matches that are still very similar to source material
- **0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

### Top Sources

- 16% Internet sources
- 14% Publications
- 8% Submitted works (Student Papers)

### Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet		
		downloads.hindawi.com	2%
2	Internet		
		jjbs.hu.edu.jo	2%
3	Internet		
		ouci.dntb.gov.ua	2%
4	Internet		
		pdfs.semanticscholar.org	1%
5	Student papers		
		University of Muhammadiyah Malang	1%
6	Publication		
		Hugo Desnos, Anne Baudot, Magda Teixeira, Gérard Louis, Loris Commin, Samuel...	1%
7	Publication		
		Maftuchah Maftuchah, Agus Zainudin, Erfan Dani Septia, Andhika Nur Sya'bani et...	1%
8	Internet		
		scholar.archive.org	1%
9	Internet		
		www.sciencepub.net	<1%
10	Internet		
		www.intechopen.com	<1%

11	Publication	Al-Musharafi S.K, Al-Amrani S.A., Mahmoud I.Y, Al-Bahry S.N, Leduc A.O.H.C. "The ...	<1%
12	Internet	sandaleromane.ro	<1%
13	Internet	www.mdpi.com	<1%
14	Publication	Dyah Roeswitawati, Robby Haikal Akbar, Zahid Hussain, Dian Indratmi, Praptinin...	<1%
15	Internet	www.bio-conferences.org	<1%
16	Publication	H. Sukorini. "Arbuscular Vascular Mycorrhizes (MVA) to Control Wilt Disease in To...	<1%
17	Internet	www.ajol.info	<1%
18	Publication	Wahyu Widodo, Adi Sutanto, Imbang Dwi Rahayu, Apriliana Devi Anggraini et al. "...	<1%
19	Publication	Yhago Patrycky Antunes Souza Assis, Anna Christina de Almeida, Wedson Carlos L...	<1%
20	Student papers	University of Technology, Sydney	<1%

# Biodiesel Industrial Waste based on *Jatropha curcas* as a Fungicide to Control *Fusarium oxysporum* and *Alternaria solani*

Henik Sukorini<sup>1\*</sup>, Dyah Erni Widyastuti<sup>1</sup>, Dini Kurniawati<sup>2</sup>, Sawita Suwannarat<sup>3</sup>, Maizirwan Mel<sup>4,5</sup>, and Roy Hendroko Setyobudi<sup>6</sup>

<sup>1</sup>Department of Agrotechnology, Faculty of Agriculture and Animal Science, University of Muhammadiyah Malang, Jl. Raya Tlogomas No. 246, Malang 65144, East Java, Indonesia

<sup>2</sup>Department of Mechanical Engineering, Faculty of Engineering, University of Muhammadiyah Malang, Malang 65144, East Java, Indonesia

<sup>3</sup>Department of Plant Pathology, Faculty of Agriculture, Kasetsart University, 50 Ngamwongwan Rd, Lat Yao, Chatuchak, Bangkok 10900, Thailand

<sup>4</sup>Department of Chemical Engineering and Sustainability, Faculty of Engineering, International Islamic University Malaysia, 50728 Kuala Lumpur, Malaysia

<sup>5</sup>Postgraduate School, Program Study of Biology, Menara Universitas Nasional, Jl. RM. Harsono No.1, Special Region of Jakarta 12550, Indonesia

<sup>6</sup>Department of Agriculture Science, Postgraduate Program, University of Muhammadiyah Malang, Malang, 65144, East Java, Indonesia

**Abstract.** *Fusarium oxysporum* (Schlecht. emend. Snyder & Hansen.) is the most devastating pathogens causing wilt disease on the tomatoes (*Solanum lycopersicum* L.) plant, whereas *Alternaria solani* (Sorauer), is a pathogen that caused early blight on potatoes (*Solanum tuberosum* L.) *Jatropha curcas* L. is a biodiesel material known as a potential fungicide. The industrial biodiesel waste based on *J. curcas* was not yet observed, particularly on the utilization of waste mainly for green manure and biogas. This research aimed to evaluate the extract of industrial biodiesel waste based on *J. curcas* against *F. oxysporum* and *A. solani*. There were  $2 \times 10^3$  mg L<sup>-1</sup>,  $4 \times 10^3$  mg L<sup>-1</sup>,  $6 \times 10^3$  mg L<sup>-1</sup>,  $8 \times 10^3$  mg L<sup>-1</sup>, and  $10 \times 10^3$  mg L<sup>-1</sup> of crude extract that were tested on both pathogens *in vitro* and *in vivo*. The concentration of the extract was in an effective range of  $10 \times 10^3$  mg L<sup>-1</sup> and was able to inhibit the growth of all isolates of *F. oxysporum* from both locations by more than 80 %, but the inhibition of the pathogen *A. solani* was less than 76 %.

**Keywords:** Environmentally friendly, pathogen control, waste to fungicide, waste utilization

## 1. Introduction

*Jatropha curcas* L. is a bioenergy resource in the family Euphorbiaceae. It is a perennial bush that is easy to grow in tropical and subtropical regions and resistant to drought. This plant is native to Mexico and Central America but can be cultivated in South East Asia [1, 2]. Indonesia has started to develop *J. curcas*-based energy to reduce dependence on fossil energy. It is possible because *J. curcas* can be grown on marginal lands that do not compete with food crops [3–6].

Tomato (*Solanum lycopersicum* L.) plants often experience decreased production due mainly to *Fusarium* wilt disease. *Fusarium* attacks tomatoes with a yield reduction of up to 85 % [3]. Until now, this disease is very difficult to control even with synthetic pesticides. Various ways have been done including using antagonistic microorganisms, but the results have not been able to effectively control this disease. The other disease is early blight. Early blight is one of the important diseases in potato plants after *Phytophthora*. The attack rate of this disease reaches 65 % [4].

In the processing of *J. curcas* as a biofuel, the process generates waste, such as seed coat waste, liquid waste, and husk. *J. curcas* seed cake is generated in considerable quantities as a by-product of *J. curcas* seed oil extraction [7, 8]. This biodiesel waste contains toxic ingredients so it cannot be used immediately. Generally, the collection and disposal

\* Corresponding author: [hsukorini@yahoo.com](mailto:hsukorini@yahoo.com)

of residues is a separate issue because the longer it is stacked, it would be problematic for the environment [9]. Burning the wastes would add more to the global CO<sub>2</sub> level resulting in the enhanced greenhouse effect and climate change [9]. On the other hand, *Jatropha* plant potentially as a fungicide, the extract of ethanol of the *Jatropha* seed showed a maximum zone of inhibitory potentials at 100 % [10], *J. curcas* seed extract showed significant antifungal activities with a growth inhibition zone of 5.6 mm or equivalent to 78.87 % inhibition followed by pulp with the zone of 7.4 mm or equivalent to 72.07 %, and whole fruits with the zone of 14.2 mm or equivalent to 46.42 % as compared to the control with the zone of 26.5 mm or equivalent to 100 % [11], as antimicrobial and insecticidal [12]. Study in utilizing biodiesel processing waste from *Jatropha* plants still have a wide opportunity, the recent study by [13–15], used two wastes of *Jatropha* that are relevant in both crop and animal production when used as organic manure and animal feeds respectively, none of the previous studies that use waste as a fungicide.

## 2 Material and methods

### 2.1 Pathogen

Pathogen *Fusarium oxysporum* (Schlecht. emend. Snyder & Hansen) causes wilt disease in tomato plants obtained from tomato production centers in Malang, Batu, and Blitar, East Java, Indonesia. Meanwhile, *Alternaria solani* (Sorauer), was isolated from the potato (*Solanum tuberosum* L.) production center of Tosari, Pasuruan, and Cangar, Batu, East Java, Indonesia. The parts of the plant which isolated were stems, leaves, roots, and soil. All isolates were pathogenic isolates. Pathogenic *F.oxysporum* and *A. solani* isolates were grown on PDA Merck for 7 d then used in a laboratory and greenhouse test.

### 2.2 Isolation of active ingredients of industrial biodiesel waste based on *Jatropha* seed

The waste of biodiesel based on *Jatropha* was obtained from the biodiesel factory. The waste taken was still fresh out of the factory. Waste extraction to obtain the active ingredients is performed by the method described by [16]. Waste of industrial biofuels from *Jatropha* seed was crushed, dried, weighed 300 g, and deep in 96 % methanol (Merck) for 3 d, then filtered by using Whatman paper No. 1. The filtrate was evaporated at 40 °C with a rotary evaporator RE-1020 at 200 mbar (1 mbar = 100 Pa). Furthermore, the dried crude extract was separated between hydrophobic and hydrophilic with dichloromethane (Merck), then the part that dissolved in dichloromethane (Merck) was evaporated and used for the antifungal test.

### 2.3 Bioassay test (*In-vitro*)

This bioassay was done to determine the inhibition ability of the crude extract to pathogens *in vitro*. Screening the best crude extract concentration were started at 0 mg L<sup>-1</sup>, 1 × 10<sup>3</sup> mg L<sup>-1</sup>, 2 × 10<sup>3</sup> mg L<sup>-1</sup>, 3 × 10<sup>3</sup> mg L<sup>-1</sup>, 4 × 10<sup>3</sup> mg L<sup>-1</sup>, 5 × 10<sup>3</sup> mg L<sup>-1</sup>, 6 × 10<sup>3</sup> mg L<sup>-1</sup>, 7 × 10<sup>3</sup> mg L<sup>-1</sup>, 8 × 10<sup>3</sup> mg L<sup>-1</sup>, 9 × 10<sup>3</sup> mg L<sup>-1</sup>, 10 × 10<sup>3</sup> mg L<sup>-1</sup>. The poisoning food method was used for screening by adding the crude extract on each concentration into a PDA medium at a Petri dish of 9 mm diameter, and then a diameter of 5 mm pathogen was placed in the center. The control treatment was by growing the pathogen on a PDA medium. All treatment was repeated 10 times. All treatments were incubated for 14 d at 27 °C, then observed for the percentage of inhibition. The formula to calculate inhibition of growth (Equation 1):

$$\text{Growth (\%)} = \frac{(\text{Colony diameter (control)} - \text{treatment})}{\text{Diameter colony on the control}} \times 100 \% \quad (1)$$

The result of the screening was used for the greenhouse test.

### 2.4 The greenhouse test

A completely randomized design was used in this test. All treatments were repeated three times. *F. oxysporum* isolates were tested on tomato plants aged 1 mo after planting, whereas isolates of *A. solani* were tested on potato plants aged 2 wk. The 100 mL isolates with a population of 10<sup>8</sup> were invested in the soil. A day after pathogen application, 100 mL of crude extract was poured into the planting medium of tomatoes or potatoes. The concentration of crude extract used in this research was the most effective concentration for inhibiting the growth of both types of pathogen in the *in vitro* test. As a positive control was a chemical pesticide, a negative control was not added to the plants. Observation variables were the incubation period and the intense attack of the disease. The incubation period was calculated from the time of inoculation until the appearance of the first symptoms of the disease. The intensity attack was calculated using the Equation 2:

$$\text{Intensity attack (\%)} = \frac{(\text{Total plant diseased})}{(\text{Total plant observed})} \times 100 \% \tag{2}$$


### Statistical analysis

The data collected were analyzed using SPSS statistical package, while means were separated using the least significant difference (LSD) test at 5 % level of significance [17, 18]


## 3 Result and discussion

### 3.1 In vitro test

The crude extracts  $10 \times 10^3 \text{ mg L}^{-1}$  of concentration were able to suppress the growth of isolates from all three origin locations of *F.oxysporum* isolates, namely the isolates from Batu (leaves) of 80.36 %, Malang (stems) of 78.46 %, Malang (roots) of 80.84 %, Blitar (stems) of 80.81 %, and Blitar leaves of 81.58 %. While the percentage of inhibition to isolate *A. solani* from Cangar (stems) of 62.2 %, Cangar (leaves) of 57.6 %, Tosari (stems and leaves) is 69.84 % and 76 % respectively (Figure 1 and Figure 2).



**Fig. 1.** The percentage of inhibition of crude extracts of biodiesel waste to the growth of various *F.oxysporum* isolates (Batu: leaf, Malang: stem, and Blitar: stem and leaf) through *in vitro* at various concentrations  $2 \times 10^3 \text{ mg L}^{-1}$ ,  $4 \times 10^3 \text{ mg L}^{-1}$ ,  $6 \times 10^3 \text{ mg L}^{-1}$ ,  $8 \times 10^3 \text{ mg L}^{-1}$ , and  $10 \times 10^3 \text{ mg L}^{-1}$



**Fig. 2** The percentage of inhibition of crude extracts of biodiesel waste to the growth of various *A. solani* isolates (stem and leaf of potatoes from Cangar and Tosari areas) through *in vitro* at various concentrations  $2 \times 10^3 \text{ mg L}^{-1}$ ,  $4 \times 10^3 \text{ mg L}^{-1}$ ,  $6 \times 10^3 \text{ mg L}^{-1}$ ,  $8 \times 10^3 \text{ mg L}^{-1}$ , and  $10 \times 10^3 \text{ mg L}^{-1}$


A concentration of  $10 \times 10^3 \text{ mg L}^{-1}$  can inhibit the growth of both types of pathogens by more than 80 %. These results indicate that the biodiesel waste made from *Jatropha* still has the potential as a fungicide as it was before the oil

was taken. In line with [19], used four extracts were acetone, methanol, ethyl acetate, aqueous, one negative control ( $0 \mu\text{L mL}^{-1}$ ), and one positive control used fungicide enriched medium (Monchamp) ( $3.33 \text{ g L}^{-1}$ ). The concentrations of various extract were  $7.5 \mu\text{L mL}^{-1}$ ,  $15 \mu\text{L mL}^{-1}$ ,  $30 \mu\text{L mL}^{-1}$ ,  $60 \mu\text{L mL}^{-1}$  and  $120 \mu\text{L mL}^{-1}$ . The radial growth inhibition test was carried out. The concentration of  $120 \mu\text{L mL}^{-1}$  had an inhibitory effect on the radial growth of the various organic extracts. Along with this result, extracts of Organic and aqueous *J. curcas* seeds could as present a fungicidal potential. Moreover, *J. curcas* leaf extracts highly inhibited mycelial growth by (85.78 %) similar to standard fungicide (Chlorothalonil) (88.37 %) in this experiment against *Phaeosariopsis personata* (Berk. & M.A. Curtis) Arx) pathogen [20].

*Jatropha* contains active ingredients that have the potential as antimicrobials. The content in castor bean extract is phenol and flavonoids which function as antimicrobial. This is consistent with the results of research from [11], in which seed extract has the potential as a natural fungicide that can replace synthetic fungicides. According to Cushnie and Lamb [21], gallic acid and pyrogallol compounds are phenolic groups that can inhibit microbial activity by acting as a chemical inhibiting the growth of microorganisms. Daidzein as isoflavonoids and routine and myricetin as flavonoids can inhibit bacterial growth by inhibiting DNA gyrase, metabolism energy, and cytoplasmic membrane. But, [20], found that among the major compounds identified with antifungal activity were hexadecanoic acid methyl ester, hexadecanoic acid ethyl ester, hexadecane, n-hexadecanoic acid, octadecanoic acid ethyl ester, phytol, and 9, 12-octadecadienoic acid (Z,Z)-methyl ester.

### 3.2 Greenhouse test

For all types of isolates, the concentration of  $10 \times 10^3 \text{ mg L}^{-1}$  may postpone the incubation period of *F.oxysporum* wilt disease more than 30 d, and it was longer than chemical pesticides by 20 d. The crude extract could delay symptom appearance of *A. solani* in more than 30 d, and it was longer than chemical pesticides (18 d). More results were presented in Figure 3 and Figure 4.




**Fig. 3.** The incubation period of *F.oxysporum* from different regions and several parts of the plant at the concentrations  $10 \times 10^3 \text{ mg L}^{-1}$


Percentage of xylem discoloration due to the attack of *F.oxysporum* origin from Batu (leaves), Blitar (Root), Blitar (leaves and stem) at a concentration of crude extract  $10 \times 10^3 \text{ mg L}^{-1}$  were respectively 4.3 cm, 4.2 cm, 4.3 cm, 4.2 cm, and 4.2 cm. It was almost the same as a positive control with chemical pesticides were an average of 4.2 cm. whereas the negative control without any application of its percentage discoloration by 9 cm (Figure 5).

Extract of biodiesel waste based on *Jatropha* was able to suppress the growth of pathogenic fungi because the extract may contain active ingredients. The presence of a phytochemical of the extract of biodiesel based on *Jatropha* can function as an antifungal [22]. On further research fractionation of industrial biodiesel extract with 80 % Me-OH using an open column chromatography could inhibit *S. rolfssii*. From this, it can be concluded that there was a secondary metabolite issued by *Jatropha* seeds that were able to inhibit the growth of *S. rolfssii* [23]. Francis [20], found that *in vitro* trial showed the antifungal activity of *J. curcas* extracts varied with the type of solvent used for extractions.

Besides, the study showed that the mycelial growth of *P. personata* was highly inhibited at the highest concentration of *J. curcas* extracts than at the lowest. This study correlated with the investigation by [24], the highest concentration (80 mg mL<sup>-1</sup>) of *J. curcas* extract inhibited the growth of *F. oxysporum* by 54 % as compared with 10 % inhibition at the lowest concentration of 20 mg mL<sup>-1</sup>. Likewise, according to [25]; disease severity was lowered as the concentration of plant extracts increased [26].



**Fig. 4.** The incubation period of *A. solani* from different regions and from several parts of the plant at at the concentrations  $10 \times 10^3$  mg L<sup>-1</sup>



**Fig. 5.** Percentage of discoloration of tomato xylem due to Fusarium wilt disease from different regions and several parts at the concentrations  $10 \times 10^3$  mg L<sup>-1</sup>

Industrial waste extract applications at a concentration of  $10 \times 10^3$  mg L<sup>-1</sup> could only delay. However, if the extract was periodically applied, as well as chemical pesticides, it was likely to be able to prevent the occurrence of these two diseases. However, when compared with the application of chemical pesticides, the extract can be an advanced alternative for environmentally-friendly disease control. Therefore, fertilizing with *jatropha*-based biodiesel production is very promising.

## 4 Conclusion

The positive results in suppressing two pathogens by crude extract of industrial biofuel waste can be an alternative for reducing chemical fungicide usage, but it is needed to be further developed and extended to make commercial fungicides.

## References

1. S.R.P. Primadari, A.K.M.A. Islam, Z. Yaakob, and S. Chakrabarty, *Jatropha curcas* L. biomass waste and its utilization, in: *Advances in biofuels and bioenergy*, M. Nageswara-Rao, and J. Soneji [eds.], (IntechOpen, London, 2018) <http://dx.doi.org/10.5772/intechopen.72803>
2. P.G. Adinurani, R.H. Setyobudi, S.K. Wahono, M. Mel, A. Nindita, E. Purbajanti, et al., *Proc. Pak. Acad. Sci.: B*, **54**, 1: 47–57 (2017) <http://ppaspk.org/index.php/PPAS-B/article/view/371>
3. M.R. Isaac, S.G. Leyva-Mr, J. Sahagun-castellanos, K. Camara-Correira, J.M. Tovar-Pedraza, J.E. Rodriguez-Perez, *Not Bot Horti Agrobot Cluj Napoca*, **46**, 2: 484–493 (2018) <https://doi.org/10.15835/nbha46211095>
4. C. Chaerani, M.K. Kardin, S. Suhardi, E. Sofiani, R.V. Glinkel, R. Groenwolt, et al., *Indones. J. Agric. Sci.* **18**: 51–62 (2018) <https://doi.org/10.21082/ijas.v18n2.2017.p51-62>
5. P.G. Adinurani, R.H. Setyobudi, S.K. Wahono, A. Sasmito, L.O. Nelwan, A. Nindita, and T. Liwang, *Int. J. Renew. Energy Dev.*, **3**, 1: 73–78 (2014) <https://doi.org/10.14710/ijred.3.1.73-78>
6. R.H. Setyobudi, T. Liwang, S. Salafudin, P.G. Adinurani, L.O. Nelwan, Y. Sakri, et al., *Energy Procedia*, **32**: 47–54 (2013) <https://doi.org/10.1016/j.egypro.2013.05.007>
7. R.H. Setyobudi, A. Wahyudi, S.K. Wahyono, P.G. Adinurani, S. Salafudin, S. Salundik, et al., *Int. J. Technol.* **4**(3): 202–208 (2013). <https://doi.org/10.14716/ijtech.v4i3.115>
8. R.H. Setyobudi, S.K. Wahono, P.G. Adinurani, S. Salafudin, A.S. Yudhanto, I. Wahyudi, et al., *Energy Procedia*, **47**: 255–262 (2014) <https://doi.org/10.1016/j.egypro.2014.01.222>
9. D. Rathore, A. Nizami, A. Singh, and D. Pant, *Biofuel Res. J.*, **3**, 2: 380–393 (2016) <https://dx.doi.org/10.18331/BRJ2016.3.2.3>
10. O.G. Cabrera, E.P.L. Molano, J. José, J.C. Álvarez, and G.A.G. Pereira, *Ceratocystis Wilt Pathogens: History and Biology—Highlighting C. cacaofunesta, the Causal Agent of Wilt Disease of Cacao*, In: *Cacao Diseases*, B. Bailey, and L. Meinhardt [eds] (Springer, Cam, 2016) [https://doi.org/10.1007/978-3-319-24789-2\\_12](https://doi.org/10.1007/978-3-319-24789-2_12)
11. M. Rahman, S.H. Ahmad, M.T.M. Mohamed, and M.Z.A Rahman, *Afr. J. Biotechnol.*, **10**, 48: 9796–9799 (2011) <https://doi.org/10.5897/AJB11.930>
12. S. Rampadarath, D. Puchooa, and R. Jeewon, *Asian Pac. J. Trop. Biomed.*, **6**, 10: 858–865 (2016) <https://doi.org/10.1016/j.apjtb.2016.01.019>
13. R.D. Akogwu, C.U. Aguorom F. Ikpa, I. Ogbonma, and J.O. Olasan, *Int. J. Environ. Agric. Biotech.*, **3**, 4: 1547–1551 (2018) <http://dx.doi.org/10.22161/ijeab/3.4.52>
14. T. Widiyastuti, T.R. Sutardi, and R.H. Setyobudi, *Energy Procedia*, **65**: 362–367 (2015) <https://doi.org/10.1016/j.egypro.2015.01.069>
15. H. Handajani, R.R. Hakim, G.A. Sutaro, B.R. Mavuso, Z.W. Chang, and S. Andriawan, *E3S Web Conf.*, **226**, 00020 (2021) <https://doi.org/10.1051/e3sconf/202122600020>
16. A.N.M. Alamgir, *Methods of Qualitative and Quantitative Analysis of Plant Constituents*. In: *Therapeutic Use of Medicinal Plants and their Extracts: Volume 2. Progress in Drug Research*, **74** (Springer, Cham, 2018) [https://doi.org/10.1007/978-3-319-92387-1\\_9](https://doi.org/10.1007/978-3-319-92387-1_9)
17. P.G. Adinurani, *Perancangan dan Analisis Data Percobaan Agro: Manual and SPSS [Design and Analysis of Agrortrial Data: Manual and SPSS]* (Plantaxia, Yogyakarta, 2016) [in Bahasa Indonesia] <https://opac.perpunas.go.id/DetailOpac.aspx?id=1159798#>
18. P.G. Adinurani, *Statistik Terapan Agroteknologi (disusun sesuai rencana pembelajaran semester) [Agrotechnology Applied Statistics (compiled according to the semester learning plan)]* (Deepublish, Yogyakarta, 2022) [in Bahasa Indonesia] <https://deepublishstore.com/shop/buku-statistika-terapan-3/>
19. N.A.N. Kone, B. Ndongo, M.M. Mountapmbeme, E.F.R. Manga, A. Heu, N.D. Mvondo, et al., *Int. J. Sci. Res. Methodol.*, **9**, 1: 95–109 (2018) [https://www.researchgate.net/profile/Francois-Manga/publication/324597606\\_Anti-Fungal\\_Activities\\_of\\_Jatropha\\_curcas\\_Seeds\\_Extracts\\_against\\_Cercospora\\_malayensis\\_Causative\\_Agent\\_of\\_Sigatoka\\_of\\_Okra\\_Leaves/links/5b11260a4585150a0a5ebdf9/Anti-Fungal-Activities-of-Jatropha-curcas-Seeds-Extracts-against-Cercospora-malayensis-Causative-Agent-of-Sigatoka-of-Okra-Leaves.pdf](https://www.researchgate.net/profile/Francois-Manga/publication/324597606_Anti-Fungal_Activities_of_Jatropha_curcas_Seeds_Extracts_against_Cercospora_malayensis_Causative_Agent_of_Sigatoka_of_Okra_Leaves/links/5b11260a4585150a0a5ebdf9/Anti-Fungal-Activities-of-Jatropha-curcas-Seeds-Extracts-against-Cercospora-malayensis-Causative-Agent-of-Sigatoka-of-Okra-Leaves.pdf)
20. M. Francis, M. Chaca, P.A. Ndakidem, and E. Mbega, *J. Anim. Plant Sci.*, **47**, 1: 8358–71 (2021) <https://dspace.nm-aist.ac.tz/handle/20.500.12479/1159>
21. T.P.T. Cushnie, and A.J. Lamb, *Int. J. Antimicrob. Agents*, **26**, 5: 343–356 (2005) <https://doi.org/10.1016/j.ijantimicag.2005.09.002>

22. S. Rachana, A. Tarun, R. Rinki, A. Neha, and R. Meghna, *J. Pharm. Biomed. Sci.*, **15**, 15: 1–4 (2012)  
[https://www.researchgate.net/profile/Rachana-Singh-4/publication/259195469\\_Comparative\\_Analysis\\_of\\_Antibacterial\\_Activity\\_of\\_Jatropha\\_curcas\\_Fruit\\_Parts/links/0c960535ddd3c0e65a000000/Comparative-Analysis-of-Antibacterial-Activity-of-Jatropha-curcas-Fruit-Parts.pdf](https://www.researchgate.net/profile/Rachana-Singh-4/publication/259195469_Comparative_Analysis_of_Antibacterial_Activity_of_Jatropha_curcas_Fruit_Parts/links/0c960535ddd3c0e65a000000/Comparative-Analysis-of-Antibacterial-Activity-of-Jatropha-curcas-Fruit-Parts.pdf)
23. S. Gopalakrishnan, I.G.K. Kannan, P. Humayun, S.V. Meesala, and D. Kanala, *Afr. J. Biotechnol.*, **9**, 47: 8048–57 (2010) <https://doi.org/10.5897/AJB10.1276>
24. M. Francis, M. Chacka, P.A. Ndakidemi, and E.R. Mbega, *Int J Agric Biol*, **26**, 2 (2021)  
<https://doi.org/10.17957/IJAB/15.1829>
25. V.K. Bajpai, K.H. Baek, E.S. Kim, J.E. Han, M.H. Kwak, K.H. Oh, et al., *Plant Pathol. J.*, **28**, 3: 317–321 (2012)  
<https://doi.org/10.5423/PPJ.NT.04.2012.0056>
26. A. Goel, and K. Sharma, *Int. J. Agric. Biol. Eng.*, **7**, 9: 859–862 (2013)  
<https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.962.1803&rep=rep1&type=pdf>