

Making Eucheuma cottonii Doty Jam with Various Palm Sugar Concentrations

by Turnitin Instructor

Submission date: 03-Nov-2023 02:13PM (UTC+0700)

Submission ID: 2215849121

File name: rniawati_2019_IOP_Conf._Ser._Earth_Environ._Sci._276_012019.pdf (758.6K)

Word count: 6045

Character count: 30637

PAPER • OPEN ACCESS

Making *Eucheuma cottonii* Doty Jam with Various Palm Sugar Concentrations

To cite this article: E C Kurniawati *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **276** 012019

View the [article online](#) for updates and enhancements.

You may also like

- [Design and Development of Stirring and Cooking Machines for Palm sugar](#)
Anwar Mazmur and Syarifuddin
- [Multi-objective optimization model for determining palm sugar granules production in remanufacturing process using NSGA-II](#)
N Umni, E Noor and M Romli
- [Nira aren's tapping business at Hasanuddin University education Forest, Maros](#)
A Mujetahid and N Dalya



245th ECS Meeting • May 26-30, 2024 • San Francisco, CA

Learn more & submit!

Present your work at the leading electrochemistry & solid-state science conference.

Network with academic, government, and industry influencers!

Submit abstracts by December 1, 2023



Making *Eucheuma cottonii* Doty Jam with Various Palm Sugar Concentrations

E C Kurniawati, Husamah*, R Latifa, S Zaenab, T I Permana, A Fauzi

Department of Biology Education, University of Muhammadiyah Malang, Jl. Raya Tlogomas No. 246 Malang 65144, East Java, Indonesia

*Corresponding author: husamahumm@gmail.com

Abstract. Seaweed is a biological resource abundantly available in Indonesia, yet it has been hardly used or consumed. This study aimed to analyze the influence of palm sugar [*Arenga pinnata* (Wumb) Merr.] concentration on the quality of *Eucheuma cottonii* Doty seaweed jam using true experimental research design. The concentrations of palm sugar were 0%, 30%, 40%, 50%, 60%, and 70%. The results showed that there was an effect of palm sugar concentration on the quality of *Eucheuma cottonii* Doty seaweed jam. The best treatment in this study was palm sugar concentration of 70%. In this treatment, water content, total sugar, and total soluble solids of seaweed jam were 33.60%, 64.89%, and 54.15%, respectively. In general, the addition of palm sugar improves seaweed jam quality.

Keywords: *Eucheuma cottonii* Doty, jam quality, palm sugar

1. Introduction

Seaweed is one of the most favored and developed marine products in all Indonesian waters and has population up to 384.73 thousand ha [1,2]. In 2010, Indonesia produced the highest amount of seaweed, harvesting 5.632 tons of wet seaweed [3]. The export of Indonesian seaweed continued to increase during the period of 2010-2014 with 14.04% export value and 11.7% export volume per annum [4]. Then, in 2016, Indonesia exported 183.257 tonnes of seaweed and earned USD 133 million from that sector [5]. So far, Indonesia and China have become the two largest seaweed producer countries [5,6]. The presence of those large quantities of seaweed originates from original stock as well as from seaweed cultivation carried out by Indonesian people [2,7-9].

Although Indonesia is positioned as the largest seaweed producer, Indonesia is not among the largest seaweed consuming countries. [5,10]. This condition is related to the unfamiliarity of the Indonesian people in consuming seaweed compared to other countries [11]. The consumption of seaweed in Indonesia is relatively low. Out of 100% total national seaweed production, 64% was exported in the form of dried seaweed while the remaining 36% was used in the domestic industry [12]. The typical aroma of seaweed seems to be the cause of the low consumption of seaweed, whether it is fresh or processed [13]. Besides, despite a large amount of seaweed in the Indonesian archipelago, sufficient effort has not been made to increase its potential.

Eucheuma cottonii Doty is a seaweed species in red algae class, which is widely cultivated in Indonesian waters [14]. *E. cottonii* is also the most dominant algae exported by Indonesia [5]. *E. cottonii* contains various essential nutrients and is useful in maintaining human's health. Some



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

bioactive compounds are also contained in these algae, such as flavonoids, hydroquinone phenols, and triterpenoids [15]. In addition, *E. cottonii* also has the potential to be a natural source of anti-tyrosinase [16]. *E. cottonii* extract also shows antioxidant activity and provides a protective effect against UV, so it is indicated to be able to be useful as a nutrient for skin rejuvenation [17]. Moreover, based on several studies, *E. cottonii* has the potential as a natural candidate for anti-cancer drugs, such as cervical cancer [18] and breast cancer [19,20].

The use of seaweed can be maximized through the diversification of seaweed products. One of the diversifications is by using *E. cottonii* as an ingredient for making jam. *E. cottonii* can be used as jam making ingredient since it contains carrageenan [21,22]. Carrageenan is a natural carbohydrate which belongs to the class of polysaccharides [23]. This carbohydrate is a hydrocolloid substance used as a thickening and gelling agents, substances needed in making jam [24,25].

In addition to the presence of hydrocolloid substances, certain conditions also need to be considered during producing seaweed jam. These conditions may affect the quality of jam and the attention of consumers. Some of these conditions, including texture, aroma, to the taste of jam produced [26,27]. In addition, the sweet taste of jam also affects the level of consumer preferences in consuming jam. The use of type and concentration of sugar will determine the sweetness and characteristics of the jam produced [28–30]. In general, sugar tables are a type of sugar used as ingredients for making jam. In fact, there are other types of sugar that have the potential as a sweetener in producing quality jam; one of them is palm sugar.

Palm sugar is a sugar commonly found in Indonesia and comes from sugar palm trees [*Arenga pinnata* (Wumb) Merr.] [31]. The tree is naturally spread in various regions in Indonesia [32–34]. Interestingly, palm sugar is also said to be healthier than table sugar [31,34]. Palm sugar is also reported to have a lower glycemic index [35]. Furthermore, besides being a natural sweetener, palm sugar also has antioxidant activities that can play a role when the body is contaminated by free radicals [36].

The utilization of *E. cottonii* as the main ingredient of jam is still rarely found. In addition, the substitution of table sugar with palm sugar in the producing jam has been seldom to be conducted. Therefore, this study will provide new information on the innovation of seaweed-based food products in Indonesia.

2. Methods

2.1. Research design

The research design used in this study was True Experimental Research to determine the differences in the concentration of palm sugar towards the quality of *E. cottonii* jam. The list of palm concentrations used in this study is presented in Table 1. The jam qualities observed in this study were from water, sugar, and total soluble solids content as well as organoleptic parameters. *E. cottonii* jam making was done at the Biology Laboratory, University of Muhammadiyah Malang (UMM), Indonesia. Water content analysis was carried out at the Animal Science and Nutrition Laboratory, UMM. Total sugar content analysis was carried out at Food Quality and Safety Testing Laboratory, Brawijaya University. The total soluble solids analysis was conducted at the Food Technology Laboratory, UMM. Finally, organoleptic tests were carried out in the Biology Laboratory, UMM.

Table1. Treatment code and palm sugar concentration

Treatment Codes*	Palm Sugar Concentration (%)
K1	0
K2	30
K3	40
K4	50
K5	60
K6	70

*code used to summarize the details of the concentration of each treatment. K = group

2.2. Materials

The main ingredients utilized in this study were *E. cottonii*. and palm sugar (*Arenga pinnata*). *E. cottonii* obtained from Sumenep, Madura, whereas the palm sugar used was "Nira Murni Gula Aren", one of palm sugar trade names from Indonesia. Seaweed used was confirmed to be *E. cottonii* Doty after the identification step was carried out at Central Laboratory of Islamic University of Malang

2.3. The production of *E. cottonii* jam

Initially, dried *E. cottonii* was cleaned from the remaining dirt. Then, the seaweed was soaked in water for three days. In each morning and afternoon, the water changed. After three days, the seaweed was dried and seaweed porridge was made by mixing 100 grams seaweed with 50 mL of water until became smooth. After that, the ingredient of seaweed jam that consisted of 200 grams seaweed, 1.5 grams' citric acid, 148.5 mL of water and palm sugar (according to each treatment in Table 1) were prepared. Those ingredients were cooked for 20 minutes at a temperature of approximately 95 °C. Finally, the seaweed jam was put into the bottles.

2.4. Water content analysis

The water content was analyzed based on the Indonesian National Standard (SNI) 2891: 1992 point 5.1. Initially, as much as 1-2 grams of sample was weighed. Then, the sample was dried in an oven at 105 °C for 3 hours. After that, the samples were cooled in a desiccator and repeated several times until the sample weight was constant. Furthermore, the water content was calculated by dividing the sample weight before drying with the sample weight after drying.

2.5. Total sugar content analysis

The testing of sugar content was done using Anthrone method. This method is divided into two major steps, making a standard curve and determining the sugar content contained in the sample. The steps for making a standard curve are as follows. Standard glucose solution of 0.0; 0.2; 0.4; 0.8, and 1.0 mL was inserted into the test tube. Then, distilled water was added to each tube until it reached 1 mL. Then, 5 mL of anthrone reagent was added to each tube. Each tube was covered with aluminum foil and homogenized. After that, each tube was heated in a water bath for 12 minutes at 100 °C. Then, the cooled solution was transferred into the cuvette and the color intensity recorded in the spectrophotometer with a wavelength of 620 nm. Then, a standard curve was created. In determining the sugar concentration from the sample, 1 mL of the sample was diluted in a test tube, and the same procedure as in making a standard curve analyzed the sample

2.6. Total soluble solids content analysis

The testing of soluble solids based on SNI 3746: 2008 used refractometer. At first, the sample was weighed as much as 40 grams and then mixed with 150 mL of water. Then, the mixture was heated to boil for 2 minutes. After that, the sample was left to stand for 20 minutes then filtered and weighed. Next, two drops of sample solution were dropped into the refractometer prism. The results of the refractometer were recorded.

2.7. Organoleptic test

Organoleptic test in this study was carried out using hedonic test method. The hedonic test is an assessment based on a person's preference level on a sample that is presented based on a hedonic scale determined by thirty assessors.

2.8. Data analysis

The data from water content, total sugar content, and total soluble solids test were analyzed using one-way ANOVA followed by Duncan's Multiple Range Test (DMRT), whereas the data from the organoleptic test were analyzed using Kruskal-Wallis H test. The prerequisite tests (normality and homogeneity test) were conducted prior to ANOVA test.

3. Results and Discussion

In this study, seaweed jam made from *E. cottonii* and palm sugar with different concentrations was produced. Seaweed jam produced from each treatment is presented in Figure 1. Based on Figure 1, each treatment could produce seaweed jam. However, the color of jam from each treatment was different. Then, the results of one-way ANOVA are summarized in Table 2. and DMRT results are presented in Table 3.



Figure 1. Seaweed jam samples produced from each treatment. The details of each treatment can be seen in Table 1

Table 2. The ANOVA Results of the effect of sugar concentration on water, sugar, and soluble solids content of jam

Sources	Df	Water Content	Sugar Content	Soluble Solids
Treatment	5	65.951*	108.768*	75.542*
Error	18			

*Significant at 5%

Table 3. The DMRT results of water, sugar, and soluble solids content of each treatment

Treatment Codes*	Water Content	Sugar Content	Soluble Solids
K1	92.80 ^e	1.56 ^a	5.42 ^a
K2	53.79 ^d	53.99 ^c	41.58 ^b
K3	50.69 ^{cd}	47.37 ^b	44.66 ^{bc}
K4	45.14 ^{bc}	45.67 ^b	48.47 ^{cd}
K5	41.50 ^b	59.27 ^{cd}	51.94 ^d
K6	33.60 ^a	64.89 ^d	54.15 ^d

Note: The scores followed by the same alphabets show that there was no significant difference on $p < 0.05$.

*The details of treatment can be seen in Table 1.

3.1. The water content of jam

Water content is the amount of water in certain materials. It usually affects the appearance, texture, and taste of food hence, it becomes one of the most important aspects in food materials [37–39]. The water content in food products like jam is also significant due to affecting its quality [40]. Based on

Table 2, it can be seen that the difference in the palm sugar concentration had a significant effect on the water content level of jam [$F(5.18) = 65.951$, $p < 0.05$]. After that, based on Table 3, K1 (palm sugar concentration of 0%) was the treatment produced jam with the highest water concentration. On the other hand, K6 (palm sugar concentration of 70%) was the treatment that had the lowest water concentration. Besides that, the data portrayed that the more sugar concentration, the lower water concentration produced. This result is in line with some previous reports, such as Khan *et al.* that studied the effect of sugar concentration on *sapodilla* jam quality [29].

Sugar contains hygroscopic molecules that can bind water [41]. The sugar molecule has a hydroxyl group which is capable of forming intramolecular hydrogen bonds. The bond produces stable hydrate and causes water to be trapped in the gel [42,43]. Because of its ability, the increase in sugar will further reduce the amount of free water.

According to SNI, the maximum water content in the jam is 35%. The K6 treatment (palm sugar concentration of 70%) produced jam with appropriate water content in accordance with SNI. Adding palm sugar concentration of less than 70% resulted in water content of jam exceeding the standard level set by SNI. The less water content obtained, the better the quality of jam produced.

The low water content in food inhibits the growth of microorganisms, so the food is not easily damaged and can last longer [44]. The low water content in food products results in the unavailability of water for the microorganism growth. This condition makes food products become more durable and long lasting because the microorganisms in the food are inactive [45]. The presence of microorganisms is the cause of food products to be easily damaged and deteriorate in quality.

3.2. Total Sugar Content in *Eucheuma cottonii* Doty seaweed jam

Total sugar content is the sugar contained in a product, which is commonly derived from sugar (sucrose), glucose, and inverted sugar (fructose and glucose) [46]. The level of sugar concentration in food products will determine the level of sweetness of the food product [47,48].

Based on Table 2, it was found that there was an effect of various concentrations of palm sugar on the total sugar content in *E. cottonii* jam [$F(5.18) = 108.768$, $p < 0.05$]. Furthermore, based on Table 3, the higher of palm sugar proportion in seaweed jam formula, the higher sugar content in seaweed jam were produced. This result is in line with some previous studies, such as Gaffar *et al.* who studied the effect of sugar concentration on orange peel jam [49].

According to SNI, the minimum sugar content in the jam is 55%. The treatment producing the right amount of total sugar content in jam as suggested by SNI was K5 treatment (palm sugar concentration of 60%) and K6 treatment (palm sugar concentration of 70%). The use of palm sugar concentration below 60% will produce jams of which total sugar content is less than the minimum level set by SNI.

3.3. Total soluble solids in *E. cottonii* jam.

The proportion of the combination of both inorganic and organic substances in a food product is commonly known as total soluble solids. Reducing sugar, non-reducing sugar, organic acids, pectin, and protein are components of the soluble solids contained in a food product [50,51]. The total value of soluble solids shows the percentage of an ingredient in a solution, which later remains as a residue resulted from evaporation and heating.

Based on the ANOVA results in Table 2, the information that can be obtained is there was a significant difference between seaweed jam that had different palm sugar concentrations in its ingredients [$F(5.18) = 75.542$, $p < 0.05$]. Then, based on DMRT results in Table 3, the highest average total soluble solids could be seen in K6 treatment (palm sugar concentration of 70%), with an average value of 54.15% mass fraction. Conversely, the lowest average total soluble solid is seen in K1 treatment (palm sugar concentration of 0%), with an average value of 5.42% mass fraction. The results of Duncan test at the significance level of 5% showed that K6 treatment (palm sugar concentration of 70%) was the best treatment, but it was not significantly different from K5 treatment (palm sugar concentration of 60%) and K4 treatment (palm sugar concentration of 50%). This result is in agreement with previous studies, such as Khan *et al.* [29].

Total soluble solids affect the viscosity of jam [26]. The higher the total soluble solids contained in a product, the more viscous the product will be [52]. Total soluble solids are increasing with the higher addition of sugar concentration during the jam making process. According to Buckle, Edward, Flet, & Wotton (2009), the total soluble solid is higher with the increasing proportion of sugar added as a result of high solubility in sugar [53]. However, the minimum total soluble solids in the jam according to SNI is 65% mass fraction. Thus, this study showed that the total soluble solid in *Eucheuma cottonii* Doty jam was less than the minimum level set by SNI.

3.4. Organoleptic results

Besides water, sugar, as well as soluble solids analysis, the organoleptic tests were also conducted on the seaweed jam produced in this study. The results of the Kruskal-Wallis H test can be seen in Table 4 and the results of the pairwise comparison is presented in Table 5. Based on Table 4, the difference in the percentage of palm sugar concentrations had a significant effect on the results of the seaweed jam's organoleptic test on color parameters [χ^2 (5) = 22.109, $p < 0.05$], aroma [χ^2 (5) = 22.181, $p < 0.05$], taste [χ^2 (5) = 22.225, $p < 0.05$], and texture [χ^2 (5) = 20.242, $p < 0.05$].

Table 4. The summary of the Kruskal-Wallis H test on the effect of palm sugar concentration on color, aroma, taste, and texture of seaweed jam

Sources	Df	Color	Aroma	Taste	Texture
Treatment	5	22.109*	22.181*	22.225*	20.242*

*Significant at 5%

Table 5. The summary of pairwise comparison through Kruskal-Wallis test adjusted with Bonferroni Correction of the six seaweeds jam produced in this study

Treatment Codes	Color	Aroma	Taste	Texture
K1	2.5a	2.5a	2.5a	2.5a
K2	8.0ab	7.0ab	8.5ab	8.75ab
K3	8.0ab	10.0ab	8.5ab	10.0ab
K4	19.5b	15.0ab	15.5ab	14.0ab
K5	18.0b	19.5b	21.5b	19.3b
K6	18.0b	21.0b	18.5b	20.5b

Note: Scores followed by the same alphabets show that there is no significant difference on $p < 0.05$. *The details of treatment can be seen in Table 1.

Table 4 shows that the highest average score of organoleptic test for color was found in K6 treatment (palm sugar concentration of 70%). On the other hand, the lowest average score was found in K1 treatment (palm sugar concentration of 0%). The jam produced in K1 treatment (palm sugar concentration of 0%) had whitish brown color. It could be from the original color of raw seaweed used in the jam ingredients. In the treatment where the concentration of palm sugar was added, seaweed jam with brown color was produced as a result of the original color of palm sugar [54].

The increasing of palm sugar concentration added to the jam during the cooking process created darker brown color in *E. cottonii* Doty seaweed jam. The darker color of the jam was resulted from the higher level of sugar concentration added to it, which was also due to the natural potential of the sugar for browning reactions, or commonly known as caramelization [55–57]. Besides that, the heating process leads to the caramelization reaction of sugar and dehydration, which forms brown color [57,58]. In addition, Buckle *et al.* (2009) believed that the brownish color of the food product was caused by caramelization of sugar resulted from high temperature and long cooking [53]. K6 treatment (palm sugar concentration of 70%) was the best and the most preferred treatment for color of *Eucheuma cottonii* Doty seaweed jam. It is seen from the higher average score of organoleptic test for color in K6 treatment compared to other treatments.

Then, Table 4 shows that palm sugar concentration of 70% treatment (K6) obtained the highest average score of aroma parameter, whereas, palm sugar concentration of 0% treatment (K1) obtained the lowest average score. K1 treatment had the lowest score due to the aroma of seaweed remained in the jam, so, most of the panelists did not like it. The seaweed aroma is one of the constraints for some people to consume either fresh or processed seaweed [13]. Organoleptic test score for the aroma of *E. Cottonii* seaweed jam tended to get higher with the increasing level of palm sugar concentration added. This case is mainly due to the effect of palm sugar aroma level. The results of the organoleptic test for aroma showed that the strong aroma of palm sugar could weaken the distinctive aroma of seaweed, which was less preferred by consumers.

In taste parameter, the highest average score of the organoleptic test was found in K6 treatment (palm sugar concentration of 70%), and the lowest average score was in K1 treatment (palm sugar concentration of 0%). The lowest score was obtained as a result of no sugar added to the jam so it did not have a sweet taste. Sugar is usually added to foods to make them more sweet [47,48,59]. In the case of *E. cottonii* Doty jam, the increasing level of palm sugar concentration added will increase the sweetness of jam produced. Usually, foods with a sweeter taste is usually more preferable [47,60,61].

In texture parameter, K6 treatment also had the highest average score, and the lowest average score was found in K1 treatment. The jam produced in K1 treatment had a soft texture, which was caused by the absence of sugar. Without sugar, the free water content rises since the water is not bound by sugar [41–43]. On the contrary, the higher concentration of palm sugar creates thick or viscous jam due to the low water content in it.

4. Conclusion

In this study, seaweed jam made from *E. cottonii* and palm sugar was produced. Several level palm sugar concentrations were used to produce the best quality of jam. The results of this study informed that there was a significant effect of palm sugar concentration level on seaweed jam quality. The best treatment was in the jam with the concentration of palm sugar as high as 70%. In this treatment, the averages of water content, sugar content, and soluble solids content were 33.60%, 64.89%, and 54.15%, respectively.

Acknowledgements

The authors would like to thank the Biology Laboratory of Universitas Muhammadiyah Malang, Indonesia, the Animal Science and Nutrition Laboratory of Universitas Muhammadiyah Malang, and the Food Quality and Safety Testing Laboratory of Brawijaya University that provided the facilities in this study. Thanks also to the Biology Education Department FTTE Universitas Muhammadiyah Malang for providing various facilities, especially research administration.

References

- [1] Kemendag 2013 Rumpun laut Indonesia *Kementrian Perdagangan Republik Indonesia*
- [2] Mulyati H and Geldermann J 2017 Managing risks in the Indonesian seaweed supply chain *Clean Technologies and Environmental Policy* **19** 175–89
- [3] Valderrama D, Cai J, Hishamunda N and Ridler N 2013 *Social and economic dimensions of carrageenan seaweed farming: a global synthesis* ed D Valderrama, J Cai, N Hishamunda and N Ridler (Rome: Food and Agriculture Organization Of The United Nations)
- [4] BPS 2015 Statistik ekspor impor Indonesia 2015
- [5] Ferdouse F, Holdt S L, Smith R, Murua P and Yang L 2018 *Global status of seaweed production* (Rome: FAO)
- [6] Buschmann A H, Camus C, Infante J, Neori A, Israel Á, Hernández-González M C, Pereda S V., Gomez-Pinchetti J L, Golberg A, Tadmor-Shalev N and Critchley A T 2017 Seaweed production: Overview of the global state of exploitation, farming and emerging research activity *European Journal of Phycology* **52** 391–406

- [7] Hendri M, Rozirwan R, Apri R and Handayani Y 2018 Intensification of seaweed cultivation *Eucheuma cottonii* with verticulture method in the water of Kelagian Island, Lampung Bay *International Journal of Marine Science* **8** 114–26
- [8] Asis A, S S S and Ilyas A 2016 Strategic policy of the government of Indonesia In the field of maritime and fisheries affairs *International Journal of Scientific and Research Publications* **6** 238–43
- [9] Eriana and Radiarta I N 2017 Observation of wild seaweed species in Labuhanbua Waters, Indonesia: A preliminary assessment for aquaculture development *OmniAkuatika* **13** 13–25
- [10] Drum R 2013 Sea vegetables for food & medicine *Well Being Journal* 3–12
- [11] Bajpai V K 2017 Korean seaweeds as a food of future: An update on use and risk factors *Indian Journal of Geo-Marine Sciences* **46** 1253–60
- [12] Kemenperin 2015 Potensi Indonesia pada olahan rumput laut *Kementrian Perindustrian*
- [13] Dwiyoitno 2011 Seaweed as a potential source of dietary fiber *Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology* **6** 9
- [14] Wijayanto T, Hendri M and Aryawati R 2011 Studi pertumbuhan rumput laut *Eucheuma cottonii* dengan berbagai metode penanaman yang berbeda di perairan Kalianda, Lampung Selatan *Maspari Journal* **03** 51–7
- [15] Nurjanah, Nurilmala M, Anwar E, Luthfiyana N and Hidayat T 2018 Identification of bioactive compounds of seaweed *Sargassum* sp. and *Eucheuma cottonii* Doty as a raw sunscreen cream *Proceedings of the Pakistan Academy of Sciences: B. Life and Environmental Sciences* vol 54 pp 311–8
- [16] Chang V S and Teo S S 2016 Evaluation of heavy metal, antioxidant and anti-tyrosinase activities of red seaweed (*Eucheuma cottonii*) *International Food Research Journal* **23** 2370–4
- [17] Lim C L, Koh R Y, Haw T Y and Boudville L A 2015 Antioxidant activity of the sea bird nest (*Eucheuma cottonii*) and its radical scavenging effect on human keratinocytes *Journal of Medical and Bioengineering* **4** 461–5
- [18] Arsianti A, Astika Y, Aziza N, Kurniasari K D, Kirana B, Mandasari D, Masita R, Zulfa F R, Dewi M K, Raisya C, Zagloel Z, Azizah N N, Putrianingsih R, Arsianti A, Astika Y, Aziza N, Masita R and Zagloel Z 2018 Phytochemical test and cytotoxic activity of macroalgae *Eucheuma cottonii* against cervical HeLa cells *Pharmacog J.* **10** 1012–7
- [19] Tan C H, Chang V S and Teo S Sen 2014 Cytotoxic activity of *Eucheuma cottonii* on MCF-7 human breast cancer *Malaysian Journal of Science* **33** 155–62
- [20] Shamsabadi F T, Khoddami A, Fard S G, Abdullah R, Othman H H and Mohamed S 2013 Comparison of tamoxifen with edible seaweed (*Eucheuma cottonii* L.) Extract in suppressing breast tumor *Nutrition and Cancer* **65** 255–62
- [21] Setha B, Mailoa M N and Gaspersz F F 2016 Analysis of quality sheet carrageenan of *Eucheuma cottonii* *International Journal of ChemTech Research* **9** 92–4
- [22] Younes M, Aggett P, Aguilar F, Crebelli R, Filipič M, Frutos M J, Galtier P, Gott D, Gundert-Remy U, Kuhnle G G, Lambré C, Leblanc J, Lillegaard I T, Moldeus P, Mortensen A, Oskarsson A, Stankovic I, Waalkens-Berendsen I, Woutersen R A, Wright M, Brimer L, Lindtner O, Mosesso P, Christodoulidou A, Ioannidou S, Lodi F and Dusemund B 2018 Re-evaluation of carrageenan (E 407) and processed *Eucheuma* seaweed (E 407a) as food additives *EFSA Journal* **16**
- [23] Necas J and Bartosikova L 2013 Carrageenan: A review *Veterinarni Medicina* **58** 187–205
- [24] Milani J and Maleki G 2012 Hydrocolloids in Food Industry *Food Industrial Processes - Methods and Equipment*
- [25] Saha D and Bhattacharya S 2010 Hydrocolloids as thickening and gelling agents in food: A critical review *Journal of Food Science and Technology* **47** 587–97
- [26] Javanmard M and Endan J 2010 A survey on rheological properties of fruit jams *International Journal of Chemical Engineering and Applications* **1** 31–7

- [27] Kayshar M S, Saifullah M, Rahman A and Uddin M B 2014 An overview of quality status of selected commercial brands of juices and jams based on public perception and laboratory analysis *J. Bangladesh Agril. Univ* **12** 183–8
- [28] Curi P N, Carvalho S, Salgado D L, Pio R, Pasqual M, Bittencourt F, Souza M De and Souza V R De 2017 Influence of different types of sugars in physalis jellies *Journal of Food Science and Technology* **37** 349–55
- [29] Khan A A, Ali S W, Manzoor S, Ayub S R and Ilyas M 2016 Influence of sugar concentration on physicochemical properties and sensory attributes of sapodilla jam *PeerJ PrePrints*
- [30] Vilela A, Matos S, Abraão A S, Lemos A M and Nunes F M 2015 Sucrose replacement by sweeteners in strawberry, raspberry, and cherry Jams: Effect on the textural characteristics and sensorial profile—A chemometric Approach *Journal of Food Processing* **2015** 1–14
- [31] Kurniawan T, Jayanudin J, Kustiningsih I and Adha Firdaus M 2018 Palm sap sources, characteristics, and utilization in Indonesia *Journal of Food and Nutrition Research* **6** 590–6
- [32] Elbersson W and Oyen L 2009 *Sugar palm (Arenga pinnata)* (FACT Foundation)
- [33] Usman A, Suman A, Hakim L and Muhaimin W 2014 The impact of home-based business processing palm augar to increase socio-economic welfare of farmers in South Halmahera Regency *IOSR Journal of Business and Management (IOSR-JBM)* **16** 32–7
- [34] Victor I R M 2015 *Processing of Arenga pinnata (Palm) sugar* (McGill University)
- [35] Srikeno K and Thongta R 2015 Effects of sugarcane, palm sugar, coconut sugar and sorbitol on starch digestibility and physicochemical properties of wheat based foods *International Food Research Journal* **22** 923–9
- [36] Winami S, Arifan F, Wisnu Broto R T D, Fuadi A and Alviche L 2018 Nira acidity and antioxidant activity of Palm sugar in Sumowono Village *Journal of Physics: Conference Series* **1025**
- [37] Molnár P J 2009 Food quality indices *Encyclopedia of Life Support Systems (EOLSS)* **II**
- [38] Miranda G, Berna À, González R and Mulet A 2015 Evolution of moisture content and texture during Storage of dried apricots *The 12th International Congress on Engineering and Food (ICEF)* (Quebec) pp 2–7
- [39] Gowen A A 2012 Water activity and food quality *Contemporary Materials* **3** 31–7
- [40] Fellows P 2011 Measuring the moisture content of foods *Food Chain* **1**
- [41] Christy A A 2014 Comparison of water adsorption characteristics of oligo and polysaccharides of α -glucose studied by near infrared spectroscopy *Advanced Materials Research* **1035** 476–82
- [42] Siringoringo M T, Sitohang A, Dewi Restuana S, Rosa T, Maruba P, Posman S, Delima P, Sisilia Y and Devi Oktavia T 2018 Effect of citric acid and sucrose concentration on the quality of passion fruit jelly with dutch eggplant *IOP Conference Series: Earth and Environmental Science* vol 205 (IOP Publishing) p 012050
- [43] Tako M 2015 The principle of polysaccharide gels *Advances in Bioscience and Biotechnology* **06** 22–36
- [44] Rawat S 2015 Food spoilage: Microorganisms and their prevention *Asian Journal of Plant Science and Research* **5** 47–56
- [45] Hamad S H 2012 Factors affecting the growth of microorganisms in food *Progress in Food Preservation* ed R Bhat, A K Alias and G Paliyath (Chichester: John Wiley & Sons) pp 405–27
- [46] Walker R W and Goran M I 2015 Laboratory determined sugar content and composition of commercial infant formulas, baby foods and common grocery items targeted to children *Nutrients* **7** 5850–67
- [47] Szulc N and Thesis B 2016 *Sugar and food additives as a part of food industry* (Centria University of Applied Sciences)

- [48] Karthik A, Shah P and Bloomfield W 2018 Deceptively sweet: Uncovering hidden sugars *Proceedings of the International Conference on Industrial Engineering and Operations Management* (Washington, DC: IEOM Society International) pp 2592–601
- [49] Gaffar R, Lahming and Rais M 2017 Pengaruh konsentrasi gula terhadap mutu selai kulit jeruk bali (*Citrus maxima*) *Jurnal Pendidikan Teknologi Pertanian* **3** 117–25
- [50] Manickavasagan A, Ganeshmoorthy K, Claereboudt M R, Al-Yahyai R and Khriji L 2014 Non-destructive measurement of total soluble solid (TSS) content of dates using near infrared (NIR) imaging *Emirates Journal of Food and Agriculture* **26** 970–6
- [51] Tasnim F, Anwar Hossain M, Nusrath S, Kamal Hossain M, Lopa D and Formuzul Haque K M 2010 Quality assessment of industrially processed fruit juices available in Dhaka City, Bangladesh *Malaysian Journal of Nutrition* **16** 431–8
- [52] Juszcak L, Witczak M, Fortuna T and Solarz B 2010 Effect of temperature and soluble solids content on the viscosity of beetroot (*Beta vulgaris*) juice concentrate *International Journal of Food Properties* **13** 1364–72
- [53] Buckle K A, Edward R A, Flet G H and Wotton M 2009 *Ilmu pangan* (Jakarta: UI Press)
- [54] Vrancken A P, Patel M K and Shen L 2014 *Sugar palm: A novel bio-ethanol feedstock: A life cycle assessment of bio-ethanol production from the Indonesian sugar palm (Arenga pinnata)* (Utrecht: Utrecht University)
- [55] Scret C 2011 *Accelerated glucose discoloration method - A quick tool for glucose stability assessment* (Lund University)
- [56] Patel K N, Modi R B, Patel H G and Apamathi K D 2013 Browning, its chemistry and implications in dairy products: A review *Indo-Am. J. Agric. & Vet. Sci.* **1** 1–12
- [57] Bastos D M, Monaro É, Siguemoto É and Séfora M 2012 Maillard reaction products in processed food: Pros and cons *Food Industrial Processes - Methods and Equipment* ed B Valdez (IntechOpen)
- [58] Golon A, Kropf C, Vockenroth I and Kuhnert N 2014 An Investigation of the complexity of maillard reaction product profiles from the thermal reaction of amino acids with sucrose using high resolution mass spectrometry *Foods* **3** 461–75
- [59] Putri E P and Fauzi A 2017 Sodium Cyclamate Effect on Nondisjunction Frequency of *Drosophila melanogaster* Meigen **8** 154–8
- [60] Hoffman A C, Salgado R V, Dresler C, Faller R W and Bartlett C 2016 Flavour preferences in youth versus adults: A review *Tobacco Control* **25** ii32–9
- [61] Appleton K, Tuorila H, Bertenshaw E, De Graaf C and Mela D 2018 Sweet taste exposure and the subsequent acceptance and preference for sweet taste in the diet: Systematic review of the published literature *American Journal of Clinical Nutrition* **107** 405–19

Making Eucheuma cottonii Doty Jam with Various Palm Sugar Concentrations

ORIGINALITY REPORT

14%

SIMILARITY INDEX

11%

INTERNET SOURCES

7%

PUBLICATIONS

2%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

1%

★ www.repo.uni-hannover.de

Internet Source

Exclude quotes On

Exclude matches Off

Exclude bibliography On