

Co-pigmentation Anthocyanins of Rose Pigment (varieties of Batu Local) with Catekin from Black Tea and Green Tea Extracts

by Mochammad Wachid

Submission date: 23-Nov-2020 12:41PM (UTC+0700)

Submission ID: 1454749114

File name: Elfi_UMM_Template_of_extended_abstrak_NP-SEA_2016_rev-1_1.pdf (109.95K)

Word count: 1132

Character count: 5993

3 Co-pigmentation Anthocyanins of Rose Pigment (varieties of Batu Local) with Catekin from Black Tea and Green Tea Extracts

Elfi Anis Saati^a, Sita Ayu Pangesti^{a*}, Sri Winarsih and Moch. Wachid^a

² ^a Food Science and Technology Department, Faculty Of Agricultural Animal Husbandry, University of Muhammadiyah Malang, East Java, Indonesia

^b Student Of Food Science and Technology Department, Faculty Of Agricultural Animal Husbandry, University of Muhammadiyah Malang East Java, Indonesia

Abstract

Anthocyanin pigment has such problems such as its sensitivity to certain conditions. Co-pigmentation can be conducted by adding substances like other pigment as catechin from a kind of tea (black and green). Methodology used in this study is randomly grouped design (RAK) method with 3 times repetition. Result of this study shows the best treatment was green tea (maceration 15 minutes) and this co-pigmentation with anthocyanin pigment extract (proportion of 1:2), contained value of soluble solid 18,88 °Brix, L (lightness) 34,81, a+ (redness) 9,20, pH 2,39, 2,61 Å (521 nm), total sugar 10,07 % and antioxidant activity 83,11 %.

Keywords: anthocyanin pigment, co-pigmentation, tea catecin

*Corresponding author: E-mail Address : elfiumm@yahoo.co.id
Telephone Number : 062563621737

1. Introduction

Anthocyanin pigment has such problems such as it is sensitive to pH (alkaline pH, > 7), temperature, sugar, oxygen, UV light and other conditions [1] that there should be treatment to make it more stable in many conditions. Co-pigmentation can be conducted by adding anion substances, kation or other substances like other pigment [2]. On preliminary study, the use of catecin can be alternative. In this study, co-pigmentation used is the best catechin from a kind of tea (black and green).

2. Methodology

This research was conducted at the Laboratory of Food Science and Technology, the Laboratory of Biotechnology and Chemistry Laboratory of the University of Muhammadiyah Malang and Chemical Laboratory of Malang State Islamic University (UIN).

This study consisted of two phases, phase -1 to determine the effect of extraction time and different types of tea used. In (the) phase 2 after have known the best catechin of both types of tea, then copigmentation with increasing concentration of catechins on the anthocyanin pigment of red roses (local of Batu, east Java) [3,4,5].

The research method in the first stage using a randomized block design (RAK) arranged as factorial, composed of two factors, the first type of tea used (black tea (A1) and green tea (A2) and the second factor was extraction duration of catechins [7] (maceration for 5 minutes (B1), 10 minutes (B2) and 15 minutes (B3). The research method used was a randomized block design with 3 times repetition.

The second of research phase is to do co-pigmentation between the best catechins with anthocyanins roses [6] on various proportions as shown below :

K0 : Control (without catechins)

K1 : The proportion of Anthocyanins catechins = 1 : 1 (v / v)

K2 : The proportion of Anthocyanins catechins = 2 : 1 (v / v)

K3 : The proportion of Anthocyanins catechins = 3 : 1 (v / v)

There are four (4) treatments that be repeated as many as three (3) times and then analyzed with a parameter of observation include total soluble solid, color intensity, pH , absorbance , the activity/antioxidant power, sugar total , the levels of catechins and anthocyanins levels.

3. Results and discussion

The difference on kind of tea and maceration duration of catecin extract do not cause interaction, yet separately the difference on kind of tea total value of solution solid (TPT), color intensity includes L value (lightness) and a+ (redness). Result of this study shows that the best treatment on stage I is A2B3 (green tea with maceration duration of 15 minutes with total solution solid (TPT) 77,1 °Brix, L value (lightness) 87,03, a+ value (redness) 1,16, b+ value (yellowness) 0,24, pH 4,64, and absorbance (240-400 nm) by value of 3,55 and antioxidant activity of 90,21 %.

Table 1. Results of Analysis of Black Tea and Green Tea as Raw Materials

Raw Material	Water Content (%)	Mineral levels (%)	Fe levels (mg / 100ml)	pH
Black tea	9,31 ± 0,05	5,59 ± 0,04	34,72 ± 0,08	5,36 ± 0,03
Green Tea	6,60 ± 0,46	5,36 ± 0,02	28,13 ± 0,16	5,82 ± 0,02

4. Conclusion

The result of this study shows that the best treatment on the first step is A2B3 (green tea with maceration duration of 15 minutes with total solution solid 77,1 °Brix, L value (lightness) 87,03, a+ value (redness) 1,16, b+ value (yellowness) 0,24, pH 4,64, and absorbance (240-400 nm) by value of 3,55 and antioxidant activity of 90,21 %.

The result of co-pigmentation between anthocyanin pigment extract and catechin extract of green tea has real influence in total solution solid, color intensity includes L,a+ and b+ values, pH, absorbance, antioxidant activity and total sugar content. The best treatment on stage II is K3 (the result of co-pigmentation between anthocyanin pigment extract and catechin green tea extract), by total value of solution solid 18,88 °Brix, L (lightness) 34,81, a+ (redness) 9,20, b+ (yellowness) 0,83, pH 2,39, absorbance (521 nm) by value of 2,61, total sugar of 10,07 %, capacity of antioxidant activity 83,11 %.

References

- [1] Li, J., 2009. *Total anthocyanin content in blue corn cookies as affected by ingredients and oven types. Dissertation.* Department of Grain Science and Industry College of Agriculture. Kansas University. Manhattan, Kansas. Pp 111
- [2] Rein, M. J. 2005. *Copigments Reaction and Color Stability of Berry Anthocyanins.* www.copigments.pdf.com.html. [31 Agustus 2009].
- [3] Saati, E.A. 2012. *Potensi Pigmen Antosianin Bunga Mawar (Rose sp) Lokal Batu Sebagai Zat Pewarna Alami Dan Komponen Bioaktif Produk Pangan.* Disertasi. Program Doktor Ilmu Pertanian. Fakultas Pertanian. Universitas Brawijaya. Malang.
- [4] _____, Simon, B.W, Yuniarta, Aulanniam. 2011. *Isolation of Red Rose Anthocyanin Pigment and Its Application to Inhibit Lipid Oxidation in Yoghurt.* *Journal of Agricultural Science and Technology B* ; 1 (2011 / 12 / 01), P95 - 98
- [5] _____, 2011 . *Isolation of Red Rose Anthocyanin Pigment and Its Application to inhibit Lipid Oxidation in Yoghurt* *Journal of Agricultural Science and Technology A 1: ISSN 1939-1250*
- [6] _____. 2014. *Invensi: Produk Pewarna Alami Makanan dari Bunga Mawar Merah (Rosa sp) Dan Proses Pembuatannya.* Granted Sertifikat No IDP000034662
- [7] Sujayanto, G. 2008. *Khasiat Teh untuk Kesehatan dan Kecantikan.* Flona Serial. Oktober (1) : hal 34-38.

Co-pigmentation Anthocyanins of Rose Pigment (varieties of Batu Local) with Catekin from Black Tea and Green Tea Extracts

ORIGINALITY REPORT

11%

SIMILARITY INDEX

11%

INTERNET SOURCES

7%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	eprints.undip.ac.id Internet Source	3%
2	Damat, Damat. "Effect of Butyrylated Arrowroot Starch to the Digesta Profile and Molar Ratio SCFA", Journal of Food Research, 2013. Publication	3%
3	sinta3.ristekdikti.go.id Internet Source	2%
4	eprints.umm.ac.id Internet Source	2%
5	web.unhas.ac.id Internet Source	1%

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Mochammad Wachid
Assignment title: Publication Articles Oktober-Desem...
Submission title: Co-pigmentation Anthocyanins of R...
File name: Elfi_UMM_Template_of_extended_...
File size: 109.95K
Page count: 2
Word count: 1,132
Character count: 5,993
Submission date: 23-Nov-2020 12:41PM (UTC+0700)
Submission ID: 1454749114

NPSEA 3RD NATURAL PIGMENTS CONFERENCE FOR SOUTH-EAST ASIA

Co-pigmentation Anthocyanins of Rose Pigment (varieties of Batu Local) with Catechin from Black Tea and Green Tea Extracts
Elfi Anis Saat¹, Sita Ayu Pangesti², Sri Winarsih and Moch. Wachid¹

¹ Food Science and Technology Department, Faculty Of Agricultural Animal Husbandry, University of Muhammadiyah Malang, East Java, Indonesia
² Student Of Food Science and Technology Department, Faculty Of Agricultural Animal Husbandry, University of Muhammadiyah Malang East Java, Indonesia

Abstract
Anthocyanin pigment has such problems such as its sensitivity to certain conditions. Co-pigmentation can be conducted by adding substances like other pigment as catechin from a kind of tea (black and green). Methodology used in this study is randomly grouped design (RAK) method with 3 times repetition. Result of this study shows the best treatment was green tea (maceration 15 minutes) and this co-pigmentation with anthocyanin pigment extract (proportion of 1:2), contained value of soluble solid 18.88 °Brix, L (lightness) 38.31, a* (redness) 9.20, pH 2.39, 2.61 Å (521 nm), total sugar 10.07 % and antioxidant activity 83.11 %.

Keywords: anthocyanin pigment, co-pigmentation, tea catechin
***Corresponding author:** E-mail Address: elfiannisa@fakho.um.ac.id
Telephone Number: 06256362137

1. Introduction
Anthocyanin pigment has such problems such as it is sensitive to pH (alkaline pH > 7), temperature, sugar, oxygen, UV light and other conditions [1] that there should be treatment to make it more stable in many conditions. Co-pigmentation can be conducted by adding anion substances, cation or other substances like other pigment [2]. On preliminary study, the use of catechin can be alternative. In this study, co-pigmentation used is the best catechin from a kind of tea (black and green).

2. Methodology
This research was conducted at the Laboratory of Food Science and Technology, the Laboratory of Biotechnology and Chemistry Laboratory of the University of Muhammadiyah Malang and Chemical Laboratory of Malang State Islamic University (UIN).
This study consisted of two phases - phase 1 to determine the effect of extraction time and different types of tea used. In (the) phase 2 after have known the best catechin of both types of tea, then copigmentation with increasing concentration of catechins on the anthocyanin pigment of red roses (local of Batu, east Java) [3-5]. The research method in the first stage using a randomized block design (RAK) arranged as factorial, composed of two factors, the first type of tea used (black tea (A1) and green tea (A2)) and the second factor was extraction duration of catechins [7] (maceration for 5 minutes (B1), 10 minutes (B2) and 15 minutes (B3). The research method used was a randomized block design with 3 times repetition.
The second of research phase is to do co-pigmentation between the best catechins with anthocyanins roses [6] on various proportions as shown below:
K1: Control (without catechins)
K2 : The proportion of Anthocyanins catechins = 1 : 1 (v / v)
K3 : The proportion of Anthocyanins catechins = 2 : 1 (v / v)
K4 : The proportion of Anthocyanins catechins = 3 : 1 (v / v)
There are four (4) treatments that be repeated as many as three (3) times and then analyzed with a parameter of observation include total soluble solid, color intensity, pH , absorbance, the activity/antioxidant power, sugar total ,the levels of catechins and anthocyanins levels.

3. Results and discussion
The difference on kind of tea and maceration duration of catechin extract do not cause interaction, yet separately the difference on kind of tea total value of solution solid (TPP), color intensity includes L, value (lightness) and a* (redness). Result of this study shows that the best treatment on stage 1 is AZB3 (green tea with maceration duration of 15 minutes with total solution solid (TPP) 77,1 °Brix, L value (lightness) 87.03, a* value (redness) 1.16, b* value (yellowness) 0.24, pH 4.64, and absorbance (240-400 nm) by value of 3.55 and antioxidant activity of 90.21 %.

Table 1. Results of Analysis of Black Tea and Green Tea as Raw Material

Raw Material	Water Content (%)	Mineral levels mg / 100ml	Brix (°Brix)	pH
Black tea	9.31 ± 0.05	5.59 ± 0.04	31.32 ± 0.08	5.36 ± 0.03
Green Tea	6.60 ± 0.06	5.36 ± 0.02	28.33 ± 0.16	5.82 ± 0.02

The 3rd Natural Pigment Conference for South-East Asia (NP-SEA) Secretariat Office
Ma Chung Research Center for Photosynthetic Pigments (MRCCPP)
Universitas Ma Chung, Villa Puncak Tidar N01, Malang 65151 East Java, INDONESIA
Tel./Fax +62 341 550 171/175, E-mail: mrccpp@machung.ac.id