


BAB 1

Beeswax formulation and Wrapping effects on red garifta mango physical characteristics.pdf

 Beeswax formulation and Wrapping effects on red garifta mango physical characteristics

 Prodi Peternakan

 University of Muhammadiyah Malang

Document Details

Submission ID

trn:oid::1:3020296594

Submission Date

Sep 25, 2024, 11:55 PM GMT+7

Download Date

Sep 25, 2024, 11:58 PM GMT+7

File Name

Beeswax_formulation_and_Wrapping_effects_on_red_garifta_mango_physical_characteristics.pdf

File Size

1.1 MB

8 Pages

2,825 Words

15,830 Characters

2% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.





Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text




Exclusions

- ▶ 11 Excluded Matches

Match Groups

-  **1 Not Cited or Quoted 2%**
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 2%  Internet sources
- 3%  Publications
- 0%  Submitted works (Student Papers)

Integrity Flags





0 Integrity Flags for Review

No suspicious text manipulations found.




Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups


-  **1** Not Cited or Quoted 2%
Matches with neither in-text citation nor quotation marks
-  **0** Missing Quotations 0%
Matches that are still very similar to source material
-  **0** Missing Citation 0%
Matches that have quotation marks, but no in-text citation
-  **0** Cited and Quoted 0%
Matches with in-text citation present, but no quotation marks

Top Sources

- 2%  Internet sources
- 3%  Publications
- 0%  Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

-  **1** Internet
sustainability.uobabylon.edu.iq 2%

PAPER • OPEN ACCESS

Beeswax formulation and Wrapping effects on red garifta mango physical characteristics

To cite this article: S Husen *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **743** 012083

View the [article online](#) for updates and enhancements.

You may also like

- [Morphological characterization and its variation of "Wirasangka" mango as the flora identity of Tegay regency, Central Java, Indonesia](#)
M Abdullah, E S Rahayu and A Retnoningsih
- [Biospeckle image processing algorithms for non-destructive differentiation between maturity and ripe stages of Indian climacteric fruits and evaluation of their ripening period](#)
S Kumari and A K Nirala
- [Comparison among different postharvest ripening treatments based on carotene contents in mango using UV-VIS and Raman spectroscopy](#)
Rahat Ullah, Saranjam Khan, Muhammad Bilal *et al.*

PRIME
PACIFIC RIM MEETING
ON ELECTROCHEMICAL
AND SOLID STATE SCIENCE

HONOLULU, HI
October 6-11, 2024

Joint International Meeting of
The Electrochemical Society of Japan (ECS)
The Korean Electrochemical Society (KECS)
The Electrochemical Society (ECS)

Early Registration Deadline:
September 3, 2024

MAKE YOUR PLANS NOW!

Beeswax formulation and Wrapping effects on red garifta mango physical characteristics

S Husen^{1*}, Muhidin¹, Sri W², Devi D S², Anggoro W¹

¹Department of Agrotechnology, Faculty of Agriculture and Animal Science, University of Muhammadiyah Malang

²Department of Food Technology, Faculty of Agriculture and Animal Science, University of Muhammadiyah Malang

Corresponding author: syarifhusen.hasan@gmail.com

Abstract. Red garifta is a mango variety that has an attractive appearance with sweet and fresh taste. It is one of the superior Indonesian fruits which having export potential. Therefore, improving the fruit quality is needed to increase market competitiveness. This study aimed to determine the effect of beeswax coating concentration and wrapping on the Red Garifta mango shelf life. There were two factors, first factor was beeswax concentration (0%, 2%, 4% and 6%), and the second factor was wrapping (without and wrapped with cardboard). The result revealed that 6% concentration beeswax is the best treatment throughout storage period with minimum loss of weight, extended color and proportional hardness. Wrapping also gave better effect on the color intensity, hardness, and less weight loss reduction. Therefore, it could be concluded that the treatment able to increase the red garifta mango shelf life.

Keywords: Fruit Waxing, Wrapping, Quality, Red Garifta

1. Introduction

Mango is tropical commodity with enormous potential for future. There are more than a thousand research about vegetative propagated mango cultivars have been published, such as Irwin (Miami), Carabo (Philippines), Alphonso (India), Bombay Green (Jamaica), Mamuang (Thailand), Tee-Vee-Dee (Ghana), and Madame Francis (Haiti) [1]. In 2010, Indonesian Tropical Fruit Research Institute also already launched new varieties called Red Garifta, which has reddish color, sweet taste, strong aroma, high moisture content (83.1-86%), sugar content (15.5" Brix), vitamin C (45.5 mg/100 g), low acid content (0.21%) and the shape is oval (14-16.5 cm) [2]. The product of mango has been developed, alike pulp, toffee, yoghurt, carbonated mango beverages, mango wine, dry mango powder, chutney, mango custard powder, mango cereal flakes, strained baby food, and mango pickle [3]. Therefore, it is necessary to develop postharvest management alike storage methods of fresh mango for enhancing its shelf life without any detrimental effect during distribution [4].

Mango is highly perishable which a very short shelf life and reach to respiration peak of ripening process. Mangoes are usually harvested at the mature-green stage, then transported to distant markets and [5] ripening in 9-12 days after harvesting period at room temperature [6] and 2-3 weeks in cold storage at 13°C [7]. This short period seriously limits the long distance commercial distribution. Several study showed that shelf-life could be extended by stored them under optimum temperature, relative humidity, atmosphere composition, and use of ripening retardants [8]. In order to extend the shelf-life



and marketing period of mango, it should be stored at 12°C and 90-95% RH, because the fruit will not suffer from chilling injury and will undergo normal ripening process [9] [10].

Waxing (edible coating) also has been used for ameliorating the postharvest fruits performance effectively and efficiency [11] [12] [13]. Moreover waxing acts as barrier on the fruits surface to suppress respiration, control moisture loss [14] [15], reducing peel dehydration, maintaining color [16], increasing titratable acidity, firmness, and effective to alleviate chilling injury [17] [18] [19] [20]. Fruit waxing had the greatest effect on preserving fruit quality, especially size, weight, and had a detrimental effect on Vitamin C [21].

Beeswax is obtained from honey combs of bees after the honey has been removed by draining or centrifuging. Beeswax consists of a complex mixture of various compounds such as hydrocarbons, esters, palmitate, palmitoleat, hidroxy palmitat and oleat esters. These vegetable oil waxes have no any noxious up shots on human health upon consumption. According to FDA, they are food grading vegetable oil waxes. This research intended to observe the effect of beeswax concentration and wrapping for enhancing mango shelf life.

2. Materials and Method

Materials used for this research were red garifta mango from CukurGondang experimental garden in Pasuruan District East Java Province Indonesia. The samples were uniform in maturity levels, it should be 80-85%, 100-105 days after the flowering period, mango weight's 315-330 gram, and free of mechanical and biological damage. The wax emulsion main ingredients were beeswax, oleic acid, water, and triethanolamine (TEA). Nested design with 1st factor : concentration of beeswax (0%, 2%, 4% dan 6%) and 2nd factor : Wrapping (without wrapped and wrapped in cardboard), with 3 replications were applied. Waxing mangoes used dipping method, which started by preparing the ingredient proportions for wax emulsion formulation (Table 1).

Table 1. Beeswax Formulation

Treatment	Beeswax (gram)	TEA (ml)	Oleic acid (ml)	Water (ml)
2%	20	7	3	970
4%	40	13	7	940
6%	60	20	10	910
8%	80	27	13	880

Beeswax was heated at temperature 90-95°C, TEA and oleic acid was added and stirred for homogenization. Wax emulsion was cooled until it reached the temperature $\pm 26^\circ\text{C}$, then it was stirred with water well until homogeneous. Red garifta mango was dipped in the wax emulsion for 10 – 30 s and dried it in room temperature, then wrapped used cardboard. The samples were taken at interval of six days for hardness, colour, and weight loss analysis.

Colour Intensity analysis

The colour skin of red garifta mango assessed by using colour reader, that expressed redness (a+) and yellowness intensity (b+).

Texture/hardness measurement

Fruit texture was assessed by using texture analyzer machine that connected with computer. The force required to penetrate 10 mm inside, using a probe 6 mm. The machine was set with compression mode at 20 mm min⁻¹, result was expressed in term of force recorded in Newton (N).

Weight loss analysis

A fruits in each block were separately marked before storage and weighed. The same fruits were consistently weighed at each sampling interval during the whole storage period. The result was expressed on a percentage basis.

$$\text{Weight loss (\%)} = \frac{W_0 - W_a}{W_0} \times 100\%$$

W_0 = weight before storage (g)

W_a = weight after storage (g)

3. Results and Discussion

3.1. Colour Intensity (Redness and Yellowness)

The results of this study indicated that beeswax concentration and wrapping were not significantly effects to the reddish and yellowish levels of red garifta mangoes (Figure 1 and 2). The color alteration of mangoes without wrapped was quicker than wrapped mangoes. The redness and yellowish intensity decreased as long as increasing of beeswax concentration.

The red mango color alteration is caused by respiration process, and the color changing speed of mangoes which stored at room temperature is faster than stored in cardboard [8]. The oxygen availability in open space more than in cardboard, so it will affect the respiration rate. The increasing beeswax concentration also affect the speed of respiration. The beeswax covered the mango peel pores, make the oxygen migration rate into the fruit is inhibited, so the rate of respiration becomes slow.




Figure 1. The redness of red garifta mango during storage

The mango ripening process involves a series of biochemical reactions, resulting into increased respiration, ethylene production, change in structural polysaccharides causing softening, degradation of chlorophyll, developing pigments by carotenoids biosynthesis, change in carbohydrates or starch conversion into sugars, organic acids, lipids, phenolics and volatile compounds, thus leading to ripening of fruit with softening of texture to acceptable quality [6] [16].




Figure 2. Yellowish intensity of red garifita mango during storage

3.2. Hardness

The ripening process of mango fruit involves a series of biochemical reactions, change in structural polysaccharides causing softening [8]. Mango fruit hardness reduced in all samples (Figure 3). Red garifita hardness without waxing and wrapped rapidly decreased during storage period, this result was consistent with experiment on Shindri var. and SufaidChaunsa var. mango [8]. The previous study treated mango by waxing beeswax 6% significantly able to retain the hardness [22]. The experiment of pineapple waxing also described that waxing was potentially useful method to alleviate chilling injury associated with the major changes of cell membrane as indicated by altering electrolyte leakage, MDA content and AsA and to maintain their quality by retarding dehydration, providing a selective barrier to moisture loss and gas exchange, which lead to reduction of general metabolism rate, improvement of textural quality and maintenance of fruit quality especially hardness[17].




Figure 3. Red garifta mango hardness during storage period

3.3. Weight loss

Fruit weight loss is mainly associated with respiration and moisture evaporation through the skin which affected by packaging [8]. The beeswax coating was effective in slowing down weight loss (Figure 4.). After stored 12 days, the unwrapped red garifta mango weight loss reached 32,82%, while waxing 6% with wrapped in cardboard reached 4,57%. The fruits with beeswax coating and kept at cardboard showed the lowest weight loss, while the highest weight loss was observed in mangoes without waxing (0%) and unwrapped.




Figure 4. Weight loss of red garifta mango during storage period

The application of wax emulsion is employed to control the weight loss, maintain its appearance and natural gloss [23]. While waxing act as barriers, thereby restricting water transfer and protecting fruit skin from mechanical injuries, as well as sealing small wounds and thus, delaying dehydration [17] [24].

- [14] Pena JE, Santos K, Baez I, Carrillo D. Physical Post-Harvest Techniques as Potential Quarantine Treatments Against *Brevipalpus yothersi* (Acarina: Tenuipalpidae). *Florida Entomol.* 2015;98(4):1169–74.
- [15] Prinsloo LC, Du Plooy W, Van Der Merwe C. Raman spectroscopic study of the epicuticular wax layer of mature mango (*Mangifera indica*) fruit. *J Raman Spectrosc.* 2004;35:561–7.
- [16] Machado FLDC, Lima RMT, Alves RE, Figueiredo RW. Influence of waxing coupled to 1-methylcyclopropene on compositional changes in early harvested ‘gold’ pineapple for export. *ActaSciAgron* [Internet]. 2014;36(2):219–25. Available from: <http://www.periodicos.uem.br/ojs/index.php/ActaSciAgron/article/view/17192>
- [17] Hu H, Li X, Dong C, Chen W. Effects of wax treatment on the physiology and cellular structure of harvested pineapple during cold storage. *J Agric Food Chem.* 2012;60:6613–9.
- [18] Meng X, Li B, Liu J, Tian S. Physiological responses and quality attributes of table grape fruit to chitosan preharvest spray and postharvest coating during storage. *Food Chem.* 2008;106:501–8.
- [19] Chien PJ, Sheu F, Lin HR. Coating citrus (*Murcotttangor*) fruit with low molecular weight chitosan increases postharvest quality and shelf life. *Food Chem.* 2007;100:1160–4.
- [20] Ahmed MJ, Singh Z, Khan AS. Postharvest Aloe vera gel-coating modulates fruit ripening and quality of “Arctic Snow” nectarine kept in ambient and cold storage. *Int J Food Sci Technol.* 2009;44(5):1024–33.
- [21] Campbell JH. Interaction of Wax, Fungicides, and Ethylene Treatments on Storage and Shelf-Life of Satsuma- Mandarins. Auburn University; 2005.
- [22] Kittur FS, Saroja N, Habibunnisa, Tharanathan RN. Polysaccharide-based composite coating formulations for shelf-life extension of fresh banana and mango. *Eur Food Res Technol.* 2001;213:306–11.
- [23] Haider STA, Ahmad S, Khan AS, Basra SMA. Comparison of different fruit coatings to enhance the shelflife of Kinnow Mandarin. *Pakistan J Agric Sci.* 2017;54(1):35–44.
- [24] Hernández-Muñoz P, Almenar E, Valle V Del, Velez D, Gavara R. Effect of chitosan coating combined with postharvest calcium treatment on strawberry (*Fragaria × ananassa*) quality during refrigerated storage. *Food Chem.* 2008;110:428–35.
- [25] Roy R, Rahim MA, Alam MS. Effect of Wrapping Papers on Physiological Changes and Shelf-life of Mango cv . Langra. *J Environ Sci Nat Resour.* 2011;4(2):99–103.