



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Artikel 5
Assignment title: Henik Sukorini 1
Submission title: Utilization of peanut shell waste and bio-slurry as organic fertil...
File name: Utilization of peanut shell waste and bio-slurry as organic fertil...
File size: 1.2M
Page count: 7
Word count: 3,052
Character count: 14,904
Submission date: 06-Aug-2025 10:25AM (UTC+0700)
Submission ID: 2725902700

IOP Conference Series:
Earth and
Environmental Science

PAPER • OPEN ACCESS

Utilization of peanut shell waste and bio-slurry as organic fertilizer for sweet corn (*Zea mays* L. Saccharata)

To cite this article: D Rosawati et al 2021 IOP Conf. Ser.: Earth Environ. Sci. 799 012047

View the [article online](#) for updates and enhancements.

PURPOSE-LED PUBLISHING

You may also like

- [Effect of Organic Matter, N, P, K, Soil, Growth and Yield of Sweet Corn in the](#)
Sugono, Ery Laitan, Anik Krisnawati et al
- [Morphological Responses of Estuarine Softshell and Sweet Corn Plants to Fertilization with Composites of Plant Compost and Three Sources of Sewer](#)
H Sigitra and Z Mulyati
- [Application of Cattle Manure Compost Mixed With Peanut Slippochloral extract as Fertilizer for Sweetcorn and Cabbage in the Field of Sweet Corn, Cabbage and Cucumber](#)
N Nurhaziq, Fikri, H Asyiani et al

UNITED THROUGH SCIENCE & TECHNOLOGY

ECS The Electrochemical Society
Advancing solid state electrochemical science & technology

Science +
Technology +
YOU!

248th
ECS Meeting
Chicago, IL
October 12-16, 2025
Hilton Chicago

Register by
September 22
to save \$\$

REGISTER NOW

This content was downloaded from IP address 202.52.52.162 on 04/08/2025 at 09:02

Artikel 5

Utilization of peanut shell waste and bio-slurry as organic fertilizer for sweet corn (*Zea mays* L. *Saccharata*)

 Henik Sukorini 1

 Publication Articles Jul - Aug 2025 Dosen UMM

 University of Muhammadiyah Malang

Document Details

Submission ID

trn:oid::1:3308949219

Submission Date

Aug 6, 2025, 10:25 AM GMT+7

Download Date

Aug 6, 2025, 11:30 AM GMT+7

File Name

Utilization of peanut shell waste and bio-slurry as organic fertilizer for sweet corn (*Zea mays* L. S....pdf

File Size

1.2 MB

7 Pages

3,052 Words

14,904 Characters

23% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.





Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text




Exclusions

- ▶ 37 Excluded Sources

Match Groups

-  **46 Not Cited or Quoted** 19%
Matches with neither in-text citation nor quotation marks
-  **8 Missing Quotations** 4%
Matches that are still very similar to source material
-  **0 Missing Citation** 0%
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted** 0%
Matches with in-text citation present, but no quotation marks

Top Sources

- 18%  Internet sources
- 20%  Publications
- 5%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- **46 Not Cited or Quoted** 19%
Matches with neither in-text citation nor quotation marks
- **8 Missing Quotations** 4%
Matches that are still very similar to source material
- **0 Missing Citation** 0%
Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted** 0%
Matches with in-text citation present, but no quotation marks

Top Sources

- 18% Internet sources
- 20% Publications
- 5% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	es.scribd.com	2%
2	Internet	earchive.tpu.ru	2%
3	Internet	www.agribalkan.net	2%
4	Publication	Herry Gusmara, Ricci Handoko Silitonga, Bilman Wilman Simanihuruk. "Growth R...	2%
5	Publication	R T Nofiyanto, V R Wati, S R Setiawati, W D Noviandi, A Kuscahyanti, E Fuskhah. " ...	1%
6	Publication	N Rumbang, Paulini, R Anjalani, E R Asie. "Application of Cattle Manure Compost ...	1%
7	Publication	Sugiono, Evy Latifah, Amik Krismawati, Sri Satya Antarlina, Listy Angraeni, Wahyu...	<1%
8	Internet	journal.jfpublisher.com	<1%
9	Internet	eprints.umm.ac.id	<1%
10	Internet	repositorio.usp.br	<1%

11	Internet	talenta.usu.ac.id	<1%
12	Internet	www.ejournal.unitaspalembang.ac.id	<1%
13	Internet	www.science.gov	<1%
14	Internet	discovery.researcher.life	<1%
15	Publication	Arthur Wallace, Richard E. Terry. "Handbook of Soil Conditioners - Substances that ...	<1%
16	Publication	Nor Hazwani Aziz, Norazwina Zainol. "Isolation and identification of soil fungi isol...	<1%
17	Internet	dichvu.cesti.gov.vn	<1%
18	Student papers	University of Huddersfield	<1%
19	Internet	digitalcommons.lsu.edu	<1%
20	Publication	F R Pambudi, A Rosmalia, I G Permana, Despal. "Fermentability and Digestibility ...	<1%
21	Internet	iaeme.com	<1%
22	Publication	Yuxuan Chen, Xiaolin Lv, Yanmin Qin, Deping Zhang et al. "Effects of different bot...	<1%
23	Internet	repository.ipb.ac.id:8080	<1%
24	Internet	garuda.kemdikbud.go.id	<1%

25	Internet	ejournal.unib.ac.id	<1%
26	Internet	pinpdf.com	<1%
27	Internet	www.mdpi.com	<1%
28	Publication	Surtinah Surtinah. "Exploring Three Varieties of Sweet Corn (Zea Mays Saccharata..."	<1%
29	Publication	G Budiyanto. "Efficacy of Organic Matter Briquette Combined with Nitrogen Fertil..."	<1%

18

PAPER • OPEN ACCESS

Utilization of peanut shell waste and bio-slurry as organic fertilizer for sweet corn (*Zea mays* L. *Saccharata*)

To cite this article: D Roeswitawati *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **759** 012047

View the [article online](#) for updates and enhancements.

You may also like

- [Application of inorganic fertilizer NK \(18-32\) on growth and yield of sweet corn on clay soil](#)
Sugiono, Evy Latifah, Amik Krismawati et al.
- [Morphological Responses of Edamame Soybean and Sweet Corn Planted in Intercropping with Combinations of Plant Populations on Three Varieties of Sweet Corn](#)
H Saputra and Z Mutaqin
- [Application of Cattle Manure Compost Mixed With Kalakai \(*Stenochlaena palustris*\) in Peatlands on Growth and Components of Yield Of Sweet Corn \(*Zea mays Saccharata*\)](#)
N Rumbang, Paulini, R Anjalani et al.

17

7

b

20

6

10

UNITED THROUGH SCIENCE & TECHNOLOGY

ECS The Electrochemical Society
Advancing solid state & electrochemical science & technology

**248th
ECS Meeting
Chicago, IL
October 12-16, 2025
Hilton Chicago**

**Science +
Technology +
YOU!**

**Register by
September 22
to save \$\$**

REGISTER NOW

3 Utilization of peanut shell waste and bio-slurry as organic fertilizer for sweet corn (*Zea mays* L. *Saccharata*)

D Roeswitawati*, M D P Arman and H Sukorini

University of Muhammadiyah Malang

*E-mail: dyahwati@umm.ac.id

Abstract. The increasing demand for sweet corn (*Zea mays*, L.) in order to meet community nutrition has encouraged farmers to make improvements to the cultivation system. One of the important factors in the process of increasing sweet corn production is soil nutrients. Peanut shell (*Arachys hypogea*, L.) and bio-slurry are organic waste, when they have undergone a decomposition process they can be used as organic fertilizers. The aim of the study was to test the correct dosage of peanut shells and the concentration of bio-slurry so that it could be used as organic fertilizer for sweet corn plants. The research was conducted in the experimental garden of the Faculty of Agriculture and Animal Husbandry, University of Muhammadiyah Malang from March to May 2020. It is a factorial experiment arranged in randomized groups, the first factor is peanut shell dose: P₁ = 10 tons / ha; P₂ = 15 tons / ha; P₃ = 20 tons / ha. The second factor is the bio-slurry concentration: B₁ = 100 ml/ L; B₂ = 125 ml/ L; B₃ = 150 ml/ L. The results showed that there was an interaction between the peanut shell dosage treatment and the bio-slurry concentration on the growth and yield of sweet corn. Dose treatment of peanut shells on growth and yield variables of sweet corn is influenced by the concentration of bio-slurry. A combination of suitable treatment for growth and yield to sweet corn is peanut shell dose 15 tons/ ha and bio-slurry concentration 125 ml/ L.

27 1. Introduction

11 Sweet corn (*Zea mays* L. *saccharata*) is a horticultural commodity favored by the community because of its taste [1, 2]. Currently, the demand for sweet corn is increasing, this encourages farmers to make improvements to the cultivation system to increase production. The production of sweet corn in East Java has increased annually, namely in 2015 amounting to 6,131,163 tonnes/ha, in 2016 amounted to 6,278,264 tonnes/ha and in 2017 amounted to 6,335,252 tonnes/ha [3, 4]. National sweet corn production has not been able to meet market demand, one of the obstacles is low productivity [5, 6]. One of the important factors for increasing sweet corn production is fertilizing and/or adding organic matter to the soil [7]. Soil organic matter is important in creating soil fertility because it can improve soil physical, chemical, and biological properties. The imbalance of the nutrient cycle in the soil is a continuing threat to the maize production system, so it is necessary to add organic sources to the soil [8]. So far, peanut shells (*Arachis hypogaea* L.) have not been fully utilized, even though peanut shells contain organic N, P, K, and C macronutrients [9]. Bio-Slurry from cow dung which is processed in a biogas reactor is a high-quality organic fertilizer which is rich in macro-nutrient content that plants need in large quantities and are essential for their growth [10, 11, 12]. The purpose of this study was to tested whether there was an interaction between the treatment of peanut skin waste dosage and bio slurry dosage treatment on the growth and yield of sweet corn, and to determine the optimal dosage of peanut and bio-slurry waste for sweet corn yield.



2 Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

2. Material and Methods

2.1. Material

The research was conducted in the experimental garden of the Faculty of Agriculture and Animal Husbandry, University of Muhammadiyah Malang from March to May 2020. It is a factorial experiment arranged in randomized groups, the first factor is peanut shell dose: P1 = 10 tons/ ha; P2 = 15 tons/ ha; P3 = 20 tons/ ha; Control = NPK fertilizer at a dose of 300 kg/ ha. The second factor is the bio-slurry concentration: B1 = 100 L/ m²; B2 = 125 L/ m²; B3 = 150 L/ m².

Slurry is taken from a cow dung disposal site in Pujon village, Pujon sub-district, Malang district. Liquid slurry is first fermented for 3 weeks or until it has no smell, then poured it onto the surface of the ground where the peanut shell has been sprinkled. Liquid slurry is fermented for 3 weeks or until it has no smell then poured onto the surface of the ground where the peanut shell has been sprinkled. The slurry dose is 400 liters / ha, while the concentration is according to treatment. Peanut shell is ground until crushed and sprinkled on the surface of the soil with a dose according to treatment.

2.2. Methods

After the soil was evenly processed, an experimental plot was made with a size of 1 m², with a distance between the plots of 50 cm. The number of experimental plots was made according to the number of combinations and was repeated three times. In one plot, planting holes were made with a spacing of 70cm x 25cm.

The crushed peanut shell waste was distributed in each plot with the dose according to treatment: P1 = 10 tons / ha; P2 = 15 ton / ha; P3 = 20 ton / ha; Control = NPK fertilizer at a dose of 300 kg / ha or 4.6 grams per plant at 7 days before planting. Likewise, the Bio-Slurry waste was sprayed on the peanut shell waste at a concentration according to the treatment: B1 = 100 ml/ L; B2 = 125 ml/ L; B3 = 150 ml/ L. The maintenance of sweet corn plants includes irrigation, weeding and planting. Watering is carried out every day until the corn plants are four weeks old. Furthermore, watering is carried out only when the plants need it. In the flowering and seed formation phases, irrigation needs to be done intensively, because in that phase the plants need more water. Routine weeding is done when the plants are one to four weeks old. After the planting is more than four weeks old, weed it only if weeds cover 50% of the plot. Heading is carried out when the plants are 3 WAP (week after planting) by filling the roots of the maize with soil. The purpose of heading is to prevent the plants from falling easily.

Observations were made on the variables: diameter of corn stem, weight of corn cob, length of corn cob, and amount dissolved solid. The data obtained were analyzed using analysis of variance, if there is a difference then it is followed by a 5% real difference test.

3. Results and Discussion

3.1. Result

The results showed that there was an interaction between the treatment of peanut shells and the concentration of bio-slurry on the growth and yield of sweet corn. After the 5% Duncan follow-up test, each treatment combination showed a significant difference. Observation variables include: stem diameter, corn cobs length, corn cobs weight, amount to dissolved solids.

3.1.1 Stem Diameter

There is an interaction between the treatment of peanut shell waste dosage and the treatment of bio-slurry waste concentration on the diameter of the corn plant. This means that every change in the dose of peanut shell waste and changes into the concentration of bio-slurry waste will have an effect on changes in the diameter of the corn plant. The 5% real difference tested results showed that there were differences between each treatment combination as presented in Figure 1.




Figure 1. Combination of doses of peanut shells and bio-slurry concentration on the diameter of maize stem.

In Figure 1, it can be seen that the P3B3 treatment combination shows the largest stem diameter followed by P3B1. While the treatment combination P1B3, P2B3, P3B2 and control did not show any significant difference. In Figure 1 it appears that the P3B3 treatment combination shows the largest stem diameter followed by P3B1. Meanwhile, the combination treatment P1B3, P2B3, P3B2 and control showed no significant difference.

3.1.2 Corn Cobs Lenght

There is an interaction between the treatment of peanut shell waste dosage and the treatment of bio-slurry waste concentration on the length of corn cobs. This means that every change in the dose of peanut shell waste and changes into the concentration of bio-slurry waste will have an effect on changes in the diameter of the corn cob lenght. The 5% significant difference test results showed that there were differences between each treatment combination as presented in Figure 2




Figure 2. Combination of doses of peanut shell and bio-slurry concentration on the length of corn cobs.

In Figure2 it can be seen that the length of corn cobs in the combination treatment of P2B3 and P3B3 does not show a significant difference based on the 5% real object test . Meanwhile , the yield of corn

cobs in the two treatment combinations was significantly different from the lengths of corn cobs in the combination of other treatments.

3.1.3 Corn Cob Weight

There is an interaction between the treatment of peanut shell waste dosage and the treatment of bio-slurry waste concentration on the weight of corn cobs. This means that every change in the dose of peanut shell waste and changes into the concentration of bio-slurry waste will have an effect on changes in the diameter of the corn cob weight. The 5% significant difference test results showed that there were differences between each treatment combination as presented in Figure 3.




Figure 3. Combination treatment of peanut shells waste dosage and bio-slurry concentration on corn cobs weight.

In Figure 3 it can be seen that the highest yield of corn cobs weight is the P3B3 combination treatment followed by the P3B1 combination treatment. Meanwhile, the combination treatment of P1B3, P2B2, and P3B2 did not show a significant difference in the yield of corn cobs weight. Likewise for the combination treatment of P1B1, P1B2, and P2B3, did not show a significant difference in the yield of corn cobs weigh.

3.1.4 Amount of Dissolved Solid

There is an interaction between the treatment of peanut shell waste dosage and the treatment of bio-slurry waste concentration on the amount of dissolved solid. This means that every change in the dose of peanut shell waste and changes into the concentration of bio-slurry waste will have an effect on changes in the diameter of the amount dissolved solid. The 5% significant difference test results showed that there were differences between each treatment combination as presented in Figure 4.




Figure 4. Combination of doses of peanut shells and bio-slurry concentration on the amount of dissolved solid.

In Figure 4, it can be seen that the dissolved solids of maize in the combination treatment of P1B2, P2B3, P3B3, and control showed no significant difference based on the results of the 5% significant difference test. This shows that the treatment of the waste user is able to produce the same amount of dissolved solids in corn as the use of inorganic NPK. However, it should be not that the use of inorganic fertilizers can damage the physical structure of the soil. On the other hand, organic waste that is returned to the soil can be used as fertilizer because it can improve the physical, chemical and biological properties of the soil so that it is beneficial to plants.

3.2. Discussion

The overall results showed that the utilization of peanut shell and bio-slurry waste was better for the growth and yield of sweet corn compared to the use of inorganic NPK fertilizers. Because organic waste is a medium for growth and reproduction for decomposing organisms such as bacteria, fungi and actinomycetes. Organic matter also improves soil chemical properties, which can increase cation exchange capacity and nutrient availability. Organic matter also improves soil biological properties, which provides a growing medium for decomposing microorganisms [13, 14, 15].

Peanut skins are waste that has not been fully utilized, even though peanut shells contain macronutrient compounds of cellulose, hemicellulose, chitin and lignin [16];[17]. Bio-slurry contains 68.59% organic matter, 17.87% C-org, 1.47% N, 0.52% P, 0.38% K, and 9.09% C / N [11]. In addition, bio-slurry contains probiotic microbes which are beneficial to the health of agricultural land and fertilize the soil [12].

The addition of organic matter in the soil at the same time can improve the physical, chemical and biological properties of the soil, so that it can be useful as an organic fertilizer [17, 18]. As the results of the study indicate that there is a synergy interaction between arbuscular mycorrhizal fungi and phosphate solubilizing bacteria, so that it can increase phosphorus uptake and plant growth [19]. Another opinion states that the presence of Zn solubilizing rhizobacteria isolated from the rhizosphere of maize plants can increase the bioavailability of Zn in the soil [20, 21]. The natural fermentation process (decomposition) of organic matter in anaerobic (without oxygen) by methane bacteria (methanogenic bacteria) so that methane gas is produced as an essential nutrient for plant growth [11].

Utilization of organic waste is a solution to the application of safe and environmentally friendly technology. Bacterial and fungal inoculums have the potential to restore fertility to degraded land through several processes. Microorganisms increase the availability of biological nutrients through nitrogen fixation and mobilization of key nutrients such as phosphorus, potassium and iron. Microbial inoculation of fungi and bacteria can help reduce the negative impact of artificial fertilizers on the environment and help restore sustainable fertility.

4. Conclusion

Base on the research results, it can be concluded as follows: there was a significant interaction between peanut shell dose treatment and bio slurry concentration on the growth and production of sweet corn. So that every change in the dosage of peanut shell waste and changes from the concentration of bi-slurry waste will affect the growth and yield of sweet corn. The combination treatment of peanut shell dose 10 ton / ha and bio slurry concentration 125 ml / L is good for variable cob corn length, the amount of dissolved solids. while the combination treatment of peanut shell dosage of 20 tons / ha and concentration of bio slurry 125 ml/ L is good for variable cob corn weight and stem diameter. Recommended combined treatments for sweet corn growth and yield are: peanut shell dose 15 tons/ ha and bio-slurry concentration 125 ml/ L.

5. References

- [1] Awalita M, Sri Darmanti, S Parman 2006 *Buletin Anatoni dan Fisiologi* **16** 20-6
- [2] Aris Satriyo W, Barunawati N, Maghfoer MD 2017 *J Produksi Tanaman* **8** 1381-1388
- [3] Badan Pusat Statistik 2018 *Ekonomi dan perdagangan. Indonesia impor jagung merdeka com* <https://www.merdeka.com>.
- [4] Umi Maryamah, SH Sutjahyo, A Hindita 2017 *Buletin Agrohorti* **5** 88-97
- [5] Meriati 2019 *J Embrio* **2** 24-35
- [6] Laksono RA, WS Nurcahyo, M Syafii 2008 *J Kultivasi* **17** 608-615
- [7] Irshad M, Ashraf M, Eneji AE, Hussain Z 2013 *J Soil Sci and Plant Nutr* **13** 115-21.
- [8] Flores D-Sanches, Pastor A, Rossing WAH, Kropp MJ, Lantinga EA. 2016 *J Soil Sci Plant Nut.* **16** 801-17
- [9] Dahlan, Darmansyah 2011. *J Agrisistem* **7** 69-78
- [10] Masi R, Cri WBY, Novaty ED 2015 *J Agrotan* **1** 45-56
- [11] Saufani I, Wawan 2017 *Jurnal Online Mahasiswa* **4** 1-11
- [12] Hilmi A, Saimul L, Tintrim R 2018 *Known Nature* **1** (1) 65-73.
- [13] Darwin HP, Ginting YC, Saputra LP, Fitri H 2017 *JHI* **8** 59-67
- [14] Precillia GP, Yunus A, Harjoko D 2018 *Agrosains* **20** 28-33
- [15] Jumini, Nurhayati, Murzani 2011 *J Floratek* **6** 165-170.
- [16] Pratiwi NE, Bistok HS, Dina B 2017 *JIP* **29** 11-20
- [17] Christina NE, Achmad A 2007 *J Sains MIPA Unila* **5** 177-82
- [18] Ramiro Gonzales M, Rinker DL 2006 *Bioresour Tech* **97** 1697-86
- [19] Shubhangi S, Company S, Ballhausen MB, Rappel S, Franken P 2020 *Microbiol Res* **240** 1-10.
- [20] Mumtaz MZ, Ahmad M, Jamil M, Hussain T 2017 *Microbiol Res* **202** 51-60
- [21] Marco A, Lampis S, Saproli, Angelini E, Vallini G 2016 *Microbiol Res* **183** 42-52

Acknowledgments

We gratefully acknowledge to University of Muhammadiyah Malang, East Java, Indonesia for supporting us to complete this research work.